arXiv:1103.0693v2 [math.SG] 26 Mar 2012

OPEN GROMOV-WITTEN INVARIANTS OF TORIC CALABI-YAU 3-FOLDS

BOHAN FANG AND CHIU-CHU MELISSA LIU

Abstract. We present a proof of the mirror conjecture of Aganagic-Vafa [3] and Aganagic-Klemm-Vafa [2] on disk enumeration in toric Calabi-Yau 3-folds for all smooth semi-projective toric Calabi-Yau 3-folds. We consider both inner and outer branes, at arbitrary framing. In particular, we recover previous results on the conjecture for (i) an inner brane at zero framing in $K_{\mathbb{P}^2}$ (Graber-Zaslow [16]), (ii) an outer brane at arbitrary framing in the resolved conifold $\mathcal{O}_{\mathbb{P}^1}(-1)\oplus\mathcal{O}_{\mathbb{P}^1}(-1)$ (Zhou [41]), and (iii) an outer brane at zero framing in $K_{\mathbb{P}^2}$ (Brini [7]).

Contents

1. Introduction 2
1.1. Open Gromov-Witten invariants 2
1.2. Large N duality: conjectures and results 3
1.3. Mirror Symmetry: conjectures and results 3
1.4. Summary of results 4
1.5. Outline of the paper 4
Acknowledgments 4
2. Geometry and Topology of Toric Calabi-Yau 3-Folds 4
2.1. Toric varieties as geometric quotients 4
2.2. Toric varieties as symplectic quotients 5
2.3. The Calabi-Yau condition 6
2.4. Aganagic-Vafa A-branes 6
2.5. Framing 6
2.6. Topology 7
3. Gromov-Witten Invariants 8
3.1. The relative FTCY 3-fold $(\hat{Y},\hat{D})$ 9
3.2. Formal relative GW invariants of $(\hat{Y},\hat{D})$ 9
3.3. Descendant GW invariants of $\hat{X}$ 11
3.4. Open GW invariants of (X,L): multiple outer branes 12
3.5. Open GW invariants of (X,L): single outer or inner brane 14
3.6. Generating functions for a fixed topological type of the domain 17
4. The Mirror Conjecture on Holomorphic Disks 18
4.1. The Hori-Vafa mirror 18
4.2. Aganagic-Vafa B-branes 19
4.3. Picard-Fuchs equations and the mirror map 19
4.4. Superpotential and the mirror prediction of disk invariants 20
5. Proof of Conjecture 4.3 22
5.1. Equivariant mirror theorem 22
5.2. Open and closed mirror maps 23
5.3. Outer brane 25
5.4. Inner brane 26

6. Explicit Mirror Formulae 27
6.1. $\mathcal{O}_{\mathbb{P}^1}(-1)\oplus\mathcal{O}_{\mathbb{P}^1}(-1)$ 27
6.2. $K_{\mathbb{P}^2}$ 29
6.3. $K_{\mathbb{P}^1\times\mathbb{P}^1}=K_{\mathbb{F}_0}$ 30
6.4. $K_{dP_1}=K_{\mathbb{F}_1}$ 31
6.5. K_{dP_2} 32
6.6. K_{dP_3} 33
6.7. The toric crepant resolution Y_m of $(\mathcal{O}_{\mathbb{P}^1}(-1)\oplus\mathcal{O}_{\mathbb{P}^1}(-1))/\mathbb{Z}_m$ 34
References 38

1. Introduction

1.1. **Open Gromov-Witten invariants.** Let X be a Kähler manifold (or more generally, an almost Kähler manifold) and let L be a Lagrangian submanifold of X. Naively, open Gromov-Witten (GW) invariants of the pair (X,L) count holomorphic maps (or more generally, J-holomorphic maps) from bordered Riemann surfaces to X with boundaries mapped into L. Open GW theory can be viewed as a mathematical theory of A-model topological open strings. String dualities have produced many striking predictions on open GW invariants, especially when X is a Calabi-Yau 3-fold and L is a special Lagrangian submanifold (or more generally, when the Maslov class of L is zero).

In this paper, we consider toric Calabi-Yau 3-folds and a particular class of Lagrangian submanifolds (called Aganagic-Vafa A-branes in this paper) introduced by Aganagic-Vafa in [3]. Let X be a toric Calabi-Yau 3-fold and L be an Aganagic-Vafa A-brane in X (in particular, L is diffeomorphic to $S^1\times\mathbb{R}^2$). The open GW invariants of the pair (X,L) depend on the following discrete data:

- topological type (g,h) of the domain bordered Riemann surface Σ, where g is the number of handles and h is the number of holes when Σ is a smooth bordered Riemann surface;
- topological type of the map $u:(\Sigma,\partial\Sigma=\coprod_{i=1}^h R_i)\to(X,L)$, which is given by the degree $\beta'=u_*[\Sigma]\in H_2(X,L;\mathbb{Z})$ and winding numbers $\mu_i=u_*[R_i]\in H_1(L;\mathbb{Z})=\mathbb{Z}$;
- framing $f\in\mathbb{Z}$ of the Aganagic-Vafa A-brane L.

More generally, L can be a disjoint union of framed A-branes $(L_1,f_1),\dots,(L_s,f_s)$, and the open GW invariants of (X,L) are

$$N^{X,L}_{g,\beta',\mu^1,\dots,\mu^s}(f_1,\dots,f_s)\in\mathbb{Q} \tag{1}$$

where $\mu^i=(\mu^i_1,\dots,\mu^i_{h_i})$ are the winding numbers associated to L_i, and $h=\sum_{i=1}^s h_i$.

When $L_1,\dots,L_s$ are outer branes, μ^i_j are positive integers, and μ^i can be viewed as a (possibly empty) partition. In this case, the authors of [29] provided a mathematical definition of the open GW invariants in (1) as formal relative GW invariants of a relative formal toric Calabi-Yau (FTCY) 3-fold $(\hat{Y},\hat{D})$[1]

$$N^{X,L}_{g,\beta',\mu^1,\dots,\mu^s}(f_1,\dots,f_s):=\sum_{\pi(\vec{d}')=\beta'}F^{\hat{Y},\hat{D}}_{g,\vec{d}',\mu^1,\dots,\mu^s}. \tag{2}$$

On the right hand side of (2), π is a surjective group homomorphism $H_2(\hat{Y};\mathbb{Z})\to H_2(X,L;\mathbb{Z})$, and the relative FTCY 3-fold $(\hat{Y},\hat{D})$ is determined by the toric Calabi-Yau 3-fold X and the *framed* Aganagic-Vafa A-branes $(L_1,f_1),\dots,(L_s,f_s)$. More details will be given in Section 3.

[1] In this paper, we use slightly different convention. See (17).

1.2. **Large N duality: conjectures and results.** The large N duality relates A-model topological string theory on Calabi-Yau 3-folds to Chern-Simons gauge theory on 3-manifolds. Motivated by the large N duality, Aganagic-Klemm-Mariño-Vafa proposed the topological vertex [1], an algorithm of computing generating functions

$$
\begin{aligned}
&F_\beta^X(\lambda) = \sum_g N_{g,\beta}^X \lambda^{2g-2},\\
&F_{\beta',\mu^1,\dots,\mu^s}^{X,L}(\lambda; f_1,\dots,f_s) = \sum_g N_{g,\beta',\mu^1,\dots,\mu^s}^{X,L}(f_1,\dots,f_s)\lambda^{2g-2+h},
\end{aligned}
\tag{3}
$$

where $\beta \in H_2(X;\mathbb{Z})$ and $N_{g,\beta}^X$ is the genus g, degree β (closed) GW invariants of X. Note that the generating functions in (3) are obtained by fixing a topological type of the map and summing over all possible topological types of the domain.

Let Z_β^X and $Z_{\beta',\mu^1,\dots,\mu^s}^{X,L}$ be the disconnected versions of F_β^X and $F_{\beta',\mu^1,\dots,\mu^s}^{X,L}$ respectively. The algorithm of AKMV consists of two steps:

- O1. Explicit gluing formula which expresses Z_β^X, $Z_{\beta',\mu^1,\dots,\mu^s}^{X,L}$ in terms of the topological vertex C_{μ^1,μ^2,μ^3}, a generating function of open GW invariants of $\mathbb{C}^3$ relative to three Aganagic-Vafa A-branes.
- O2. Evaluation of the topological vertex C_{μ^1,μ^2,μ^3} by relating it to the colored HOMFLY polynomial of a 3-component link in S^3.

In [29], Li-Liu-Liu-Zhou developed a mathematical theory of the topological vertex using algebraic relative GW theory developed by J. Li [26, 27][2]:

- R0. Open GW invariants of (X, L) are defined as formal relative GW invariants of a relative FTCY 3-fold $(\hat{Y}, \hat{D})$.
- R1. The degeneration formula satisfied by these formal relative GW invariants agrees with the gluing formula in [1], with C_{μ^1,μ^2,μ^3} replaced by a generating function $\widetilde{C}_{\mu^1,\mu^2,\mu^3}$ of invariants of an in-decomposable relative FTCY 3-fold.
- R2. LLLZ evaluated $\widetilde{C}_{\mu^1,\mu^2,\mu^3}$ using virtual localization and a formula of Hodge integrals provided in [28], and showed that it agrees with the formula of C_{μ^1,μ^2,μ^3} in [1] when one of the partitions is empty.

This proves the topological vertex algorithm up to 2-leg vertex. The validity of the full 3-leg case is a consequence of the proof of Gromov-Witten/Donaldson-Thomas correspondence of toric Calabi-Yau 3-folds by Maulik-Okounkov-Oblomkov-Pandharipande [35].

1.3. **Mirror Symmetry: conjectures and results.** The mirror symmetry relates A-model topological string theory on a Calabi-Yau 3-fold X to the B-model topological string theory on the mirror Calabi-Yau 3-fold $X^\vee$.

Let X be a smooth toric Calabi-Yau 3-fold. We assume that X is *semi-projective*, i.e., X has a torus fixed point and X is projective over its affinization $\mathrm{Spec}\big(H^0(X,\mathcal{O}_X)\big)$. By the results in [21, Section 2], a smooth toric variety is semi-projective if and only if it is equal to the GIT quotient of an affine space $\mathbb{C}^r$ by the action of a subtorus of $(\mathbb{C}^*)^r$. Let L be an Aganagic-Vafa A-brane in X. Aganagic-Vafa related a generating function of disk invariants of the pair (X, L) to the Abel-Jacobi map of the mirror curve of X [3]. It was clarified in [2] that the framing of L corresponds to choice of flat coordinates in the B-model. The integrals in [3, 2] are solutions to extended Picard-Fuchs equations [23, 24, 36, 37]. To our knowledge, the above mirror conjectures on disk invariants has been verified in the following cases: (i) X is the total space of $K_{\mathbb{P}^2}$ and L is an inner brane at zero framing (T. Graber and E. Zaslow [16]); (ii)X is the resolved conifold (i.e. the total space of

[2]The symplectic relative GW theory was developed independently by Li-Ruan [25] and Ionel-Parker [18, 19].

$\mathcal{O}_{\mathbb{P}^1}(-1)\oplus\mathcal{O}_{\mathbb{P}^1}(-1))$ and L is an outer brane at arbitrary framing (J. Zhou [41]); (iii) X is the total space of $K_{\mathbb{P}^2}$ and L is an outer brane at zero framing (A. Brini [7, Section 5.3]).

Based on the work of Eynard-Orantin [10] and Mariño [34], Bouchard-Klemm-Mariño-Pasquetti [5] proposed a new formalism of the B-model topological strings on the mirrors of toric Calabi-Yau 3-folds, and provided an recursive algorithm, now known as the remodeling conjecture, which determines the generating functions

$$
\begin{aligned}
&F_g^X(Q)=\sum_{\beta\in H_2(X;\mathbb{Z})} N_{g,\beta}^X Q^\beta\\
&F_{g,h}^{X,L}(Q,x_1,\dots,x_h;f)=\sum_{\substack{\beta'\in H_2(X,L;\mathbb{Z})\\ \mu=(\mu_1,\dots,\mu_h)}} N_{\beta',\mu_1,\dots,\mu_h}(f)Q^{\beta'}x_1^{\mu_1}\cdots x_h^{\mu_h}
\end{aligned}
\tag{4}
$$

from the disk invariants $F_{0,1}^{X,L}$ and annulus invariants $F_{0,2}^{X,L}$. Note that the generating functions are obtained by fixing the topological type of the domain and summing over all possible topological types of the map. The remodeling conjecture has been proved for the framed 1-leg topological vertex by L. Chen [8] and by J. Zhou [39]. J. Zhou later proved the conjecture for the framed 3-leg topological vertex [40]. Recently, Bouchard-Catuneanu-Marchal-Sułkowski proved the remodeling conjecture for $F_g^{\mathbb{C}^3}$ [4].

1.4. **Summary of results.** The goals of the present paper are twofold.

(i) We use localization to derive a formula of formal relative GW invariants of the relative FTCY 3-fold $(\hat{Y},\hat{D})$ in terms of descendant GW invariants of the FTCY 3-fold $\hat{X}:=\hat{Y}-\hat{D}$ (**Proposition 3.1**). As a consequence, we obtain a formula of open GW invariants of (X,L) in terms of descendant GW invariants of X when L is a union of framed outer branes (**Proposition 3.2**). The formula allows us to extend the definition to a single framed inner brane (**Proposition 3.4**).

(ii) We use the approach in Graber-Zaslow [16] to prove the mirror conjecture on disk invariants in [3, 2, 23, 24, 36, 37] for any smooth semi-projective toric Calabi-Yau 3-folds. We consider both outer and inner branes, at arbitrary framing (**Conjecture 4.3**). We list explicit mirror formulae for the resolved conifold, local toric Fano surfaces, and toric crepant resolutions of $\big(\mathcal{O}_{\mathbb{P}^1}(-1)\oplus\mathcal{O}_{\mathbb{P}^1}(-1)\big)/\mathbb{Z}_m$.

1.5. **Outline of the paper.** In Section 2, we review the geometry and topology of toric Calabi-Yau 3-folds and Aganagic-Vafa A-branes. In Section 3, we introduce and compare various GW invariants: formal relative GW invariants of $(\hat{Y},\hat{D})$, descendant GW invariants of $\hat{X}$, open GW invariants of (X,L), and descendant GW invariants of X. In Section 4, we review the geometry of mirrors of toric Calabi-Yau 3-folds and Aganagic-Vafa B-branes, and state the mirror conjecture on holomorphic disks (Conjecture 4.3). The proof of Conjecture 4.3 is given in Section 5. We list explicit mirror formulae in Section 6.

Acknowledgments. We thank C. Woodward, E. Zaslow, and J. Zhou for their comments.

2. Geometry and Topology of Toric Calabi-Yau 3-Folds

2.1. **Toric varieties as geometric quotients.** We refer to [12] for the theory of toric varieties. In this subsection, we consider smooth, possibly noncompact toric varieties of any dimension.

Let $N\cong\mathbb{Z}^n$ be a free abelian group, and let $\triangle$ be a *fan* in N (or in $N_{\mathbb{R}}=N\otimes\mathbb{R}\cong\mathbb{R}^n$) of strongly convex rational polyhedral cones. Let $X=X(\triangle)$ be the toric variety associated to $\triangle$. In this paper, we assume that X is smooth. We use the following notation:

- $\triangle(d)$ is the set of d-dimensional cones in $\triangle$.

- Given $\sigma \in \triangle(d)$, let $V(\sigma)$ denote the codimension d orbit closure associated to σ.
- Let $\triangle(1) = \{\rho_1, \ldots, \rho_r\}$ be the set of 1-dimensional cones in $\triangle$, and let $v_i \in N$ be the unique generator of the semigroup $\rho_i \cap N$, so that $\rho_i \cap N = \mathbb{Z}_{\geq 0} v_i$.
- Let $M = \mathrm{Hom}(M, \mathbb{Z})$ be the dual lattice of N.

There is a group homomorphism

$$\phi : \widetilde{N} := \bigoplus_{i=1}^{r} \mathbb{Z}\widetilde{v}_i \cong \mathbb{Z}^r \longrightarrow N \cong \mathbb{Z}^n, \quad \widetilde{v}_i \mapsto v_i.$$

We assume that ϕ is surjective. Let

$$l^{(a)} = (l_1^{(a)}, \ldots, l_r^{(a)}), \quad a = 1, \ldots, k,$$

be a $\mathbb{Z}$-basis of $\mathrm{Ker}\phi \cong \mathbb{Z}^k$, where $k = r - n$. Let $\widetilde{M} = \mathrm{Hom}(M; \mathbb{Z})$ be the dual lattice of $\widetilde{N}$. Since $\phi : \widetilde{N} \to N$ is surjective, the dual map $\phi^* : M \to \widetilde{M}$ is injective. Applying $\mathrm{Hom}(-, \mathbb{C}^*)$ to the following short exact sequence

$$0 \to M \xrightarrow{\phi^*} \widetilde{M} \to A_{n-1}(X) \to 0,$$

we obtain another short exact sequence

$$1 \to G \to \widetilde{\mathbb{T}} \to \mathbb{T} \to 1, \tag{5}$$

where

$$\begin{aligned} G &:= \mathrm{Hom}(A_{n-1}(X), \mathbb{C}^*) \cong (\mathbb{C}^*)^k, \\ \widetilde{\mathbb{T}} &= \mathrm{Hom}(\widetilde{M}, \mathbb{C}^*) \cong (\mathbb{C}^*)^r, \\ \mathbb{T} &= \mathrm{Hom}(M, \mathbb{C}^*) \cong (\mathbb{C}^*)^n. \end{aligned}$$

We also denote

$$\mathbb{L} = \mathrm{Ker}\phi, \quad \mathbb{L}^\vee = A_{n-1}(X).$$

Notice that $\mathbb{L}^\vee \cong H^2(X; \mathbb{Z})$, and $\mathbb{L} \cong H_2(X; \mathbb{Z})$.

The torus $\widetilde{\mathbb{T}} \cong (\mathbb{C}^*)^r$ acts on $\mathbb{C}^r = \mathrm{Spec}\mathbb{C}[X_1, \ldots, X_r]$. Let $I(\triangle) \subset \mathbb{C}[X_1, \ldots, X_r]$ be the ideal generated by $\{\prod_{\rho_i \not\subset \sigma} X_i \mid \sigma \in \triangle\}$, and let $Z(\triangle) \subset \mathbb{C}^r$ be the subvariety defined by $I(\triangle)$. Then X can be identified with the geometric quotient:

$$X = (\mathbb{C}^r - Z(\triangle))/G. \tag{6}$$

where $G = (\mathbb{C}^*)^k$ acts on $\mathbb{C}^r$ by

$$(t_1, \ldots, t_k) \cdot (X_1, \ldots, X_r) = (\prod_{a=1}^{k} t_a^{l_1^{(a)}} \cdot X_1, \ldots, \prod_{a=1}^{k} t_a^{l_r^{(a)}} \cdot X_r).$$

The $\widetilde{\mathbb{T}}$-divisor $\{X_i = 0\}$ in $\mathbb{C}^r - Z(\triangle)$ descends to a $\mathbb{T}$-divisor D_i in X. When X is semi-projective, the quotient (6) is also a quotient in the sense of GIT (geometric invariant theory) [21, Section 2].

2.2. **Toric varieties as symplectic quotients.** When X is a smooth semi-projective toric variety, we may describe Kähler structures on X as a symplectic quotient. Let $G_{\mathbb{R}} \cong U(1)^k$ be the maximal compact subgroup of $G \cong (\mathbb{C}^*)^k$. Then the dual $\mathfrak{g}_{\mathbb{R}}^*$ of the Lie algebra $\mathfrak{g}_{\mathbb{R}}$ of $G_{\mathbb{R}}$ can be canonically identified with $A_{n-1}(X) \otimes \mathbb{R} = H^{1,1}(X_\Sigma; \mathbb{R})$. Let $\widetilde{\mu} : \mathbb{C}^r \to \mathfrak{g}_{\mathbb{R}}^* \cong \mathbb{R}^k$ be the moment map of the Hamiltonian $G_{\mathbb{R}}$-action on $\mathbb{C}^r$, equipped with the standard Kähler form

$$\frac{\sqrt{-1}}{2} \sum_{i=1}^{r} dX_i \wedge d\overline{X}_i. \tag{7}$$

Then

$$\widetilde{\mu}(X_1,\dots,X_r)=(\sum_{i=1}^r l_i^{(1)}|X_i|^2,\dots,\sum_{i=1}^r l_i^{(k)}|X_i|^2).$$

Let $(r_1,\dots,r_k)\in H^{1,1}(X;\mathbb{R})\cong\mathbb{R}^k$ be a Kähler class. Then

$$X=\widetilde{\mu}^{-1}(r_1,\dots,r_k)/G_{\mathbb{R}}. \tag{8}$$

The standard Kähler form (7) on $\mathbb{C}^r$ descends to a Kähler form $\omega_{r_1,\dots,r_k}$ on the quotient (8) in class $(r_1,\dots,r_k)\in H^{1,1}(X;\mathbb{R})$. The real numbers $r_1,\dots,r_k$ are known as *Kähler parameters* of X. Let $T_a=-r_a+\sqrt{-1}\theta_a$ be *complexified Kähler parameters* of X.

2.3. **The Calabi-Yau condition.** We say a toric variety X is Calabi-Yau if the canonical divisor $K_X=-D_1-\cdots-D_r$ is trivial. X is Calabi-Yau if and only if

$$\sum_{i=1}^r l_i^{(a)}=0,\quad a=1,\dots,k$$

When X is Calabi-Yau, we have a short exact sequence

$$1\to\mathbb{T}'\to\mathbb{T}\to\mathbb{C}^*\to 1, \tag{9}$$

where $\mathbb{T}'\cong(\mathbb{C}^*)^{n-1}$ is the subtorus of $\mathbb{T}$ that acts trivially on $\Lambda^n T_X\cong\mathcal{O}_X$.

2.4. **Aganagic-Vafa A-branes.** Let $X=\widetilde{\mu}^{-1}(r_1,\dots,r_k)/G_{\mathbb{R}}$ be a smooth semi-projective toric Calabi-Yau 3-fold equipped with the Kähler form $\omega_{r_1,\dots,r_k}$. Here $\widetilde{\mu}^{-1}(r_1,\dots,r_k)$ is defined by

$$\sum_{i=1}^{k+3} l_i^{(a)}|X_i|^2=r_a,\quad a=1,\dots,k.$$

Write $X_i=\rho_i e^{\sqrt{-1}\phi_i}$, where $\rho_i=|X_i|$. In [3], Aganagic-Vafa introduced Lagrangian submanifolds L of X_Σ defined by additional constraints:

$$\sum_{i=1}^{k+3}\hat{l}_i^1|X_i|^2=c_1,\quad \sum_{i=1}^{k+3}\hat{l}_i^2|X_i|^2=c_2,\quad \sum_{i=1}^{k+3}\phi_i=\text{const},$$

where

$$\hat{l}_i^\alpha\in\mathbb{Z},\quad \sum_{i=1}^{k+3}\hat{l}_i^\alpha=0,\quad \alpha=1,2.$$

Such a Lagrangian submanifold is diffeomorphic to $S^1\times\mathbb{C}$ and intersects a unique 1-dimensional orbit closure $V(\tau_L)$, where $\tau_L\in\triangle(2)$, along a circle. We say L is an *outer* brane if $V(\tau_L)\cong\mathbb{C}$, and we say L is an *inner* brane if $V(\tau_L)\cong\mathbb{P}^1$.

2.5. **Framing.** Let $\mathbb{T}'_{\mathbb{R}}\cong U(1)^2$ be the maximal compact subgroup of $\mathbb{T}'\cong(\mathbb{C}^*)^2$. Then $\mathbb{T}'_{\mathbb{R}}$ preserves any Aganagic-Vafa A-brane L. The $\mathbb{T}'_{\mathbb{R}}$-action on L is given by

$$(t_1,t_2)\cdot(z,u)=(t_1z,t_2u)$$

where $(t_1,t_2)\in\mathbb{T}'_{\mathbb{R}}=U(1)\times U(1)$, $(z,u)\in L\cong S^1\times\mathbb{C}$. Therefore (X,L) is a $\mathbb{T}'_{\mathbb{R}}$-equivariant pair. The $\mathbb{T}'_{\mathbb{R}}$-action on X is Hamiltonian with respect to the symplectic form $\omega_{r_1,\dots,r_k}$. Let $\mu':X\to(\mathfrak{t}'_{\mathbb{R}})^*$ be the moment map of the $\mathbb{T}'_{\mathbb{R}}$-action, where $(\mathfrak{t}'_{\mathbb{R}})^*\cong\mathbb{R}^2$ is the dual of the Lie algebra $\mathfrak{t}'_{\mathbb{R}}$ of $\mathbb{T}'_{\mathbb{R}}$.

We define the 1-skeleton of X to be

$$X^1=\bigcup_{\tau\in\triangle(2)}V(\tau).$$

Then $\mu'(X^1)$ is an immersed trivalent graph. $\mu'(V(\tau))$ is a line segment if $V(\tau) \cong \mathbb{P}^1$, and is a ray if $V(\tau) \cong \mathbb{C}$. Suppose that σ is a 3-dimensional cone in $\triangle$ and $\tau_L \subset \sigma$. Then $V(\sigma) \subset V(\tau_L)$, where $V(\sigma)$ is the $\mathbb{T}$-fixed point associated to σ. Let

$$\mathsf{u}_L = (c_1)_{\mathbb{T}'}(T_{V(\sigma)}V(\tau_L)) \in H^2_{\mathbb{T}'}(V(\sigma);\mathbb{Z}) \cong H^2(B\mathbb{T}';\mathbb{Z}) \subset H^2(B\mathbb{T}';\mathbb{R}) \cong (\mathfrak{t}'_{\mathbb{R}})^*.$$

Then u_L is tangent to $\mu(V(\tau_L))$. Let $\tau'_L \in \triangle(2)$ be the 2-cone such that $\tau'_L \subset \sigma$ and $\mu'(V(\tau'_L))$ is the first edge encountered in the counterclockwise direction from $\mu'(V(\tau_L))$. Define

$$\mathsf{v}_L = (c_1)_{\mathbb{T}'}(T_{V(\sigma)}V(\tau'_L)) \in H^2_{\mathbb{T}'}(V(\sigma);\mathbb{Z}) \subset (\mathfrak{t}'_{\mathbb{R}})^*.$$

Let $\rho_{i_1} \subset \sigma$ be the 1-cone that is not in τ_L, and let $\rho_{i_2} \subset \sigma$ be the 1-cone that is not in τ'_L. We use ρ_{i_3} to denote the 1-cone in σ other than ρ_{i_1}, ρ_{i_2}. If $V(\tau_L) \cong \mathbb{P}^1$, then there is another 3-cone $\sigma' \in \triangle(3)$ other than σ of which τ_L is a face. Let ρ_{i_4} be the 1-cone in σ' but not in τ_L.



FIGURE 1. images of $D_{i_1}, D_{i_2}, D_{i_3}, D_{i_4}$, $V(\sigma)$, $V(\sigma')$ under μ'.

Let L be an Aganagic-Vafa A-brane. Then $\mu'(L)$ is a point. A framing of L is a choice of a vector $\mathsf{f} \in H^2(B\mathbb{T}';\mathbb{Z})$ such that $\mathsf{u}_L \wedge \mathsf{f} = \mathsf{u}_L \wedge \mathsf{v}_L$, i.e., $\mathsf{f} = \mathsf{v}_L - f\mathsf{u}_L$ for some $f \in \mathbb{Z}$. We sometimes call the integer f the a framing of L. See Figure 2.



FIGURE 2. framing of an A-brane

We may view f as an element in $\mathrm{Hom}(\mathbb{T}', \mathbb{C}^*)$. Let $\mathbb{T}_{L,f} \cong \mathbb{C}^*$ be the kernel of $\mathsf{f} : \mathbb{T}' \to \mathbb{C}^*$.

2.6. **Topology.** Suppose that $p_1, \ldots, p_k$ is a $\mathbb{Z}$-basis of $H^2(X;\mathbb{Z}) \cong \mathbb{L}^\vee$ dual to $l^{(1)}, \ldots, l^{(k)} \in \mathbb{L}$. We choose appropriate $\{l^{(a)}\}_{a=1}^k$ and $\{p_a\}_{a=1}^k$ such that the class of $\omega_{r_1,\ldots,r_k}$ is $\sum_{a=1}^k r_a p_a \in H^2(X_\Sigma;\mathbb{R}) = H^{1,1}(X_\Sigma;\mathbb{R})$. Let $Q_a = e^{T_a}$, which are formal Kähler parameters. Given $\beta \in H_2(X;\mathbb{Z})$, define

$$Q^\beta := e^{\langle \beta, \sum_{a=1}^k T_a p_a \rangle} = \prod_{a=1}^k Q_a^{d_a},$$

where $d_a = \langle \beta, p_a \rangle$, $a = 1, \ldots, k$. For $i = 1, \ldots, r$, we have

$$D_i^* := c_1(\mathcal{O}_X(D_i)) = \sum_{a=1}^k l_i^{(a)} p_a, \quad \langle D_i^*, \beta \rangle = \sum_{a=1}^k d_a l_i^{(a)}.$$

Let $\triangle(2)_c = \{\tau \in \triangle(2) \mid V(\tau) \cong \mathbb{P}^1\}$. The inclusion $X^1 \hookrightarrow X$ induces a surjective group homomorphism

$$H_2(X^1;\mathbb{Z}) = \bigoplus_{\tau \in \triangle(2)_c} V(\tau) \to H_2(X;\mathbb{Z}).$$

Let

$$L = L_1 \cup \cdots \cup L_s$$

be a disjoint union of s Aganagic-Vafa A-branes $L_1, \ldots, L_s$ in X. We have the following long exact sequence of relative homology groups of the pair (X, L):

$$\cdots \to H_2(L;\mathbb{Z}) \to H_2(X;\mathbb{Z}) \to H_2(X,L;\mathbb{Z}) \xrightarrow{\partial} H_1(L;\mathbb{Z}) \to H_1(X;\mathbb{Z}) \to \cdots \tag{10}$$

where $H_2(L;\mathbb{Z}) = 0$ and $H_1(X;\mathbb{Z}) = 0$.

For $i = 1, \ldots, s$, let $V(\tau_i)$ be the unique 1-dimensional orbit closure that intersects L_i. We assume that $\tau_1, \ldots, \tau_s$ are distinct. We choose a $\mathbb{T}$-fixed point $x_i \in V(\tau_i)$. When L_i is an outer brane, $V(\tau_i) \cong \mathbb{C}$, and there is a unique such point; when L_i is an inner brane, there are exactly two such points, x_i and x_i^-. Then $V(\tau_i) - L_i \cap V(\tau_i)$ has two connected components, one of which is a holomorphic disk B_i containing x_i. We orient B_i by the holomorphic structure. Then B_i represents a relative homology class $b_i \in H_2(X, L;\mathbb{Z})$. Let $\gamma_i \in H_1(L_i;\mathbb{Z})$ be the image of b_i under the map ∂ in (10). Then

$$H_1(L;\mathbb{Z}) = \bigoplus_{i=1}^{s} H_1(L_i;\mathbb{Z}) = \bigoplus_{i=1}^{s} \mathbb{Z}\gamma_i.$$

Any element $\beta' \in H_2(X, L;\mathbb{Z})$ is of the form

$$\beta' = \beta + \sum_{i=1}^{s} d_i b_i,$$

where $d_i \in \mathbb{Z}$ and $\beta \in H_2(X;\mathbb{Z})$.

Let $\mathcal{A}$ be the collection of *anti-cones*

$$\mathcal{A} = \{J \subset \{1, \ldots, r\} | \sum_{j \notin J} \mathbb{R}_{\geq 0} v_j \text{ is a cone in } \Delta\}.$$

The effective cone $\mathbb{L}_{\text{eff}}$ is

$$\mathbb{L}_{\text{eff}} = \{\beta \in \mathbb{L} | \{i \in \{1, \ldots, r\} | \langle D_i^*, \beta\rangle \in \mathbb{Z}_{\geq 0}\} \in \mathcal{A}\}.$$

Usually the cone $\mathbb{L}_{\text{eff}} \otimes_{\mathbb{Z}} \mathbb{R}$ in the real vector space $\mathbb{L} \otimes_{\mathbb{Z}} \mathbb{R}$ is called the Mori cone. One says $\beta \geq 0$ if $\beta \in \mathbb{L}_{\text{eff}}$ and $\beta > 0$ if β is also non-zero. The closure of the Kähler cone in $\mathbb{L}^\vee \otimes_{\mathbb{Z}} \mathbb{R}$ is

$$C_X = \bigcap_{I \in \mathcal{A}} (\sum_{i \in I} \mathbb{R}_{\geq 0} v_i).$$

It is dual to the Mori cone.

3. Gromov-Witten Invariants

Let X be a toric Calabi-Yau 3-fold defined by a fan $\triangle$. Let

$$(L_1, f_1), \ldots, (L_s, f_s)$$

be framed Aganagic-Vafa A-branes (see Section 2.4), and let $\tau_1, \ldots, \tau_s \in \triangle(2)$ be defined as in Section 2.6. Let $\sigma_1, \ldots, \sigma_s \in \triangle(3)$ be the top dimensional cones corresponding to $\mathbb{T}$ fixed points $x_1, \ldots, x_s$, respectively, where x_i are defined as in Section 2.6. Then

$$\tau_i \subset \sigma_i, \quad V(\sigma_i) = \{x_i\}, \quad i = 1, \ldots, s.$$

Define

$$w_1^i = \mathsf{u}_{L_i}, \quad w_2^i = \mathsf{v}_{L_i}, \quad w_3^i = -w_1^i - w_2^i, \quad \mathsf{f}_i = w_2^i - f_i w_1^i.$$

In Section 3.1–3.4, we assume $L_1, \ldots, L_s$ are outer branes. In Section 3.5 –3.6, $L = L_1$ is a single outer or inner brane.

3.1. **The relative FTCY 3-fold** $(\hat{Y}, \hat{D})$**.** We refer to [29, Section 3] for the definition of formal toric Calabi-Yau (FTCY) graphs and construction of relative FTCY 3-folds. Let Γ_X be the formal toric Calabi-Yau (FTCY) graph associated to the smooth toric 3-fold X (see [29, Section 3.1]). We define a FTCY graph

$$\Gamma_{X,(L_1,f_1),\ldots,(L_s,f_s)}$$

by replacing the noncompact edge $\bar{e}_i$ in Γ_X associated to τ_i by a compact edge $\bar{e}'_i$ with framing f_i.[3] Figure 3.1 shows the construction near the edge $\bar{e}_i$.



FIGURE 3. The FTCY graphs Γ_X and $\Gamma_{X,(L_1,f_1),\ldots,(L_s,f_s)}$. The trivalent vertex v_i corresponds to the cone $\sigma_i \in \triangle(3)$.

Let $(\hat{Y}, \hat{D})$ be the relative FTCY 3-fold associated to the FTCY graph $\Gamma = \Gamma_{X,(L_1,f_1),\ldots,(L_s,f_s)}$. Then $\hat{D}$ is the disjoint union of its connected components $\hat{D}^1, \ldots, \hat{D}^s$ which correspond to the s univalent vertices $v'_1, \ldots, v'_s$ in Γ. $\mathbb{T} = (\mathbb{C}^*)^3$ and $\mathbb{T}' = (\mathbb{C}^*)^2$ acts on $(\hat{Y}, \hat{D})$. Let $E_c(\Gamma_X)$ and $E_c(\Gamma)$ denote the set of compact edges in Γ_X and Γ, respectively. Then $E_c(\Gamma) = E_c(\Gamma_X) \cup \{\bar{e}'_1, \ldots, \bar{e}'_s\}$. Each compact edge $\bar{e}$ corresponds to a torus invariant projective line $C^{\bar{e}}$ in X or in $\hat{Y}$. We have

$$\begin{aligned} H_2(X^1;\mathbb{Z}) &= \bigoplus_{\bar{e}\in E_c(\Gamma_X)} \mathbb{Z}[C^{\bar{e}}], \\ H_2(\hat{Y};\mathbb{Z}) &= \bigoplus_{\bar{e}\in E_c(\Gamma)} \mathbb{Z}[C^{\bar{e}}] = H_2(X^1;\mathbb{Z}) \oplus \bigoplus_{i=1}^{s} \mathbb{Z}[C^{\bar{e}_i}], \end{aligned}$$

where $C^{\bar{e}_i} = V(\tau_i)$. There is a surjective group homomorphism

$$\pi : H_2(\hat{Y};\mathbb{Z}) = H_2(X^1;\mathbb{Z}) \oplus \bigoplus_{i=1}^{s} \mathbb{Z}[C^{\bar{e}_i}] \longrightarrow H_2(X, L;\mathbb{Z}) = H_2(X;\mathbb{Z}) \oplus \bigoplus_{i=1}^{s} \mathbb{Z}b_i$$

given by

$$\beta + \sum_{i=1}^{s} d_i[C^{\bar{e}_i}] \mapsto i_*\beta + \sum_{i=1}^{s} d_i b_i,$$

where $\beta \in H_2(X^1;\mathbb{Z})$, and $i_* : H_2(X^1;\mathbb{Z}) \to H_2(X;\mathbb{Z})$ is induced by the inclusion $i : X^1 \hookrightarrow X$.

3.2. **Formal relative GW invariants of** $(\hat{Y}, \hat{D})$**.** We refer to [29, Section 4] for the precise definitions of relative stable morphisms to a relative FTCY 3-fold, and formal relative GW invariants. Here we briefly recall the definitions for our particular setting.

An *effective class* of Γ is a pair $(\vec{d}, \vec{\mu})$, where

- $\vec{d} : E_c(\Gamma) \to \mathbb{Z}_{\geq 0}$ is a map from the set of compact edges in Γ to the set of nonnegative integers,
- $\vec{\mu} : \{v'_1, \ldots, v'_s\} \to \mathcal{P}$ is a map from the set of univalent vertices in Γ to set of partitions,

[3] In [29] all the noncompact edges are replaced by a compact edge ending at an univalent vertex. Here we only compactify $\bar{e}_1, \ldots, \bar{e}_s$.

- $\vec{\mu}(v_i')$ is a partition of $\vec{d}'(\bar{e}_i')$.

$\vec{\mu}$ can be viewed as an s-uple of partitions: $\vec{\mu}=(\mu^1,\dots,\mu^s)$, where $\mu^i:=\vec{\mu}(v_i')$.

Define

$$\ell(\vec{\mu})=\sum_{i=1}^{s}\ell(\mu^i),\quad \mathrm{Aut}(\vec{\mu})=\prod_{i=1}^{s}\mathrm{Aut}(\mu^i).$$

We also use $\vec{d}'$ to denote the following effective curve class:

$$\sum_{\bar{e}\in\Gamma}\vec{d}'(\bar{e})[C^{\bar{e}}]\in H_2(\hat{Y};\mathbb{Z}).$$

A genus g, class $(\vec{d}',\vec{\mu})$ relative stable morphism to $(\hat{Y},\hat{D})$ is an $\ell(\vec{\mu})$-pointed, genus g, degree $\vec{d}'$ stable morphism $u:(C,\mathbf{q})\to\hat{Y}$, where

$$\mathbf{q}=\{q_j^i\mid 1\le i\le s, 1\le j\le \ell(\mu^i)\}$$

are distinct smooth points on C, and

$$u^{-1}(\hat{D}^i)=\sum_{i=1}\mu_j^i q_j^i$$

as Cartier divisors. Let $\mathcal{M}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})$ be the moduli space of genus g, class $(\vec{d}',\vec{\mu})$ relative stable morphisms to $(\hat{Y},\hat{D})$. The stable compactification $\overline{\mathcal{M}}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})$ of $\mathcal{M}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})$ contains relative morphisms to expanded targets $(\hat{Y}_{\mathbf{m}},\hat{D}_{\mathbf{m}})$. Then $\overline{\mathcal{M}}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})$ is a formal Deligne-Mumford stacks equipped with a perfect obstruction theory of virtual dimension $\ell(\vec{\mu})$. Moreover, $\mathbb{T}'=(\mathbb{C}^*)^2$ acts $\overline{\mathcal{M}}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})$, and the perfect obstruction theory are $\mathbb{T}'$-equivariant. The genus g, class $(\vec{d}',\vec{\mu})$ formal relative GW invariants of $(\hat{Y},\hat{D})$ is defined by

$$F^{\hat{Y},\hat{D}}_{g,\vec{d}',\vec{\mu}}=\frac{1}{|\mathrm{Aut}(\vec{\mu})|}\int_{[\overline{\mathcal{M}}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})^{\mathbb{T}'}]^{\mathrm{vir}}}\frac{\prod_{i=1}^{s}\prod_{j=1}^{\ell(\mu^i)}(\mathrm{ev}_j^i)^*(c_1)_{\mathbb{T}'}(\mathcal{O}_{\hat{D}_i}(\hat{L}_i))}{e_{\mathbb{T}'}(N^{\mathrm{vir}})},\tag{11}$$

where $\overline{\mathcal{M}}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})^{\mathbb{T}'}$ is the $\mathbb{T}'$ fixed points substack in $\overline{\mathcal{M}}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})$, ev_j^i is the evaluation at the marked point q_j^i, $\hat{L}^i$ is a $\mathbb{T}'$-equivariant divisor in $\hat{D}^i$, and N^{vir} is the virtual normal bundle of $\overline{\mathcal{M}}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})^{\mathbb{T}'}$ in $\overline{\mathcal{M}}_{g,\vec{d}',\vec{\mu}}(\hat{Y},\hat{D})$. Let $\{\mathsf{u},\mathsf{v}\}$ be a $\mathbb{Z}$-basis of $H^2(B\mathbb{T}';\mathbb{Z})$, so that $H^*(B\mathbb{T}';\mathbb{Q})=\mathbb{Q}[\mathsf{u},\mathsf{v}]$. A priori the right hand side of (11) is a rational function in u,v, homogeneous of degree 0, but it is indeed a rational number independent of u,v.

Introduce variables $\mathbf{p}=\{p_j^i\mid 1\le i\le s, j\in\mathbb{Z}_{>0}\}$ and $Q=\{Q_{\bar{e}}\mid \bar{e}\in E_c(\Gamma)\}$. We define generating functions of formal relative GW invariants of $(\hat{Y},\hat{D})$:

$$F^{\hat{Y},\hat{D}}_{\vec{d}',\vec{\mu}}(\lambda)=\sum_g N^{\hat{Y},\hat{D}}_{\vec{d}',\vec{\mu}}\lambda^{2g-2+\ell(\vec{\mu})},\tag{12}$$

$$F^{\hat{Y},\hat{D}}(\lambda,Q,\mathbf{p})=\sum_{\substack{(\vec{d}',\vec{\mu})\\ \vec{\mu}\neq(\emptyset,\dots,\emptyset)}}F^{\hat{Y},\hat{D}}_{\vec{d}',\vec{\mu}}(\lambda)Q^{\vec{d}'}\mathbf{p}_{\vec{\mu}}\tag{13}$$

where $Q^{\vec{d}'}=\prod_{\bar{e}\in E_c(\Gamma)}Q_{\bar{e}}^{\vec{d}'(\bar{e})}$, $\mathbf{p}_{\vec{\mu}}=\prod_{i=1}^{s}\prod_{j=1}^{\ell(\mu^i)}p^i_{\mu^i_j}$. Define generating functions $F^{\bullet\hat{Y}\hat{D}}_{\vec{d}',\vec{\mu}}$ and $Z^{\hat{Y},\hat{D}}$ by

$$Z^{\hat{Y},\hat{D}}(\lambda,Q,\mathbf{p})=\exp\left(F^{\hat{Y},\hat{D}}(\lambda,q,\mathbf{p})\right)=\sum_{(\vec{d}',\vec{\mu})}F^{\bullet\hat{Y},\hat{D}}_{\vec{d}',\vec{\mu}}(\lambda)Q^{\vec{d}'}\mathbf{p}_{\vec{\mu}}.\tag{14}$$

3.3. **Descendant GW invariants of $\hat{X}$.** Let $\hat{X}$ be the formal completion of X along the 1-skeleton X^1. Then $\hat{X} = \hat{Y} - \hat{D}$ is a FTCY 3-fold, and

$$H_2(\hat{X};\mathbb{Z}) = H_2(X^1;\mathbb{Z}) = \bigoplus_{\bar{e}\in E_c(\Gamma_X)} \mathbb{Z}[C^{\bar{e}}].$$

The surjective group homomorphism $\pi : H_2(\hat{Y};\mathbb{Z}) \to H_2(X,L;\mathbb{Z})$ restricts to a surjective group homomorphism $H_2(\hat{X};\mathbb{Z}) \to H_2(X;\mathbb{Z})$.

An effective class of Γ_X is a function $\vec{d} : E_c(\Gamma_X) \to \mathbb{Z}_{\geq 0}$; we also use $\vec{d}$ to denote the effective curve class

$$\sum_{\bar{e}\in E_c(\Gamma_X)} \vec{d}(\bar{e})[C^{\bar{e}}] \in H_2(\hat{X};\mathbb{Z}).$$

Let $\overline{\mathcal{M}}_{g,n}(\hat{X},\vec{d})$ be the moduli space of n-pointed, genus g, degree $\vec{d}$ stable morphisms to $\hat{X}$. Define

$$\begin{aligned} G^{\hat{X}}_{g,\vec{d},\vec{\mu}}(\mathsf{u},\mathsf{v}) =& \frac{(-\sqrt{-1})^{\ell(\vec{\mu})}}{|\mathrm{Aut}(\vec{\mu})|} \prod_{i=1}^{s}\prod_{j=1}^{\ell(\mu^i)} \frac{\prod_{m=1}^{\mu^i_j-1}(\mu^i_j \mathsf{w}^i_2 + m\mathsf{w}^i_1)}{\mu^i_j \cdot \mu^i_j!(\mathsf{w}^i_1)^{\mu^i_j-1}} \cdot \\ & \int_{[\overline{\mathcal{M}}_{g,\ell(\vec{\mu})}(\hat{X},\vec{d})^{\mathbb{T}'}]^{\mathrm{vir}}} \frac{1}{e_{\mathbb{T}'}(N^{\mathrm{vir}})} \prod_{i=1}^{s}\prod_{j=1}^{\ell(\mu^i)} \frac{(\mathrm{ev}^i_j)^*\phi_i}{\frac{\mathsf{w}^i_1}{\mu^i_j}(\frac{\mathsf{w}^i_1}{\mu^i_j} - \psi^i_j)} \end{aligned} \tag{15}$$

where $\phi_i \in H_{\mathbb{T}'}(\hat{X};\mathbb{Z})$ is the $\mathbb{T}'$-equivariant Poincaré dual of the torus fixed point $x_i \in V(\tau_i)$, and

$$\mathrm{ev}^i_j : \overline{\mathcal{M}}_{g,\ell(\vec{\mu})}(\hat{X},\vec{d}) \to \hat{X}, \quad i = 1,\ldots,s, \quad j = 1,\ldots,\ell(\mu^i),$$

are evaluations at the marked points.

Define generating functions

$$\begin{aligned} G^{\hat{X}}_{\vec{d},\vec{\mu}}(\lambda;\mathsf{u},\mathsf{v}) &= \sum_g \lambda^{2g-2+\ell(\vec{\mu})} G^{\hat{X}}_{\vec{d},\vec{\mu}}(\mathsf{u},\mathsf{v}) \\ G^{\hat{X}}(\lambda,Q,\mathbf{p};\mathsf{u},\mathsf{v}) &= \sum_{\substack{(\vec{d},\vec{\mu}) \\ \vec{\mu}\neq(\emptyset,\ldots,\emptyset)}} G^{\hat{X}}_{\vec{d},\vec{\mu}}(\lambda;\mathsf{u},\mathsf{v}) Q^{\vec{d}} \mathbf{p}_{\vec{\mu}} \\ G^{\bullet\hat{X}}(\lambda,Q,\mathbf{p};\mathsf{u},\mathsf{v}) &= \exp\Big(G^{\hat{X}}(\lambda,Q,\mathbf{p};\mathsf{u},\mathsf{v})\Big) = \sum_{(\vec{d},\vec{\mu})} G^{\bullet\hat{X}}_{\vec{d},\vec{\mu}}(\lambda;\mathsf{u},\mathsf{v}) Q^{\vec{d}} \mathbf{p}_{\vec{\mu}} \end{aligned}$$

Given two partitions μ^+, μ^- of d, define

$$\Phi^{\bullet}_{\mu^+,\mu^-}(\lambda) = \sum_{\nu\vdash d} e^{\kappa_\nu \lambda/2} \frac{\chi_\nu(\mu^+)}{z_{\mu^+}} \frac{\chi_\nu(\mu^-)}{z_{\mu^-}}$$

where $\kappa_\mu = \sum_{j=1}^{\ell(\mu)} \mu_j(\mu_j - 2j + 1)$, $z_\mu = \mathrm{Aut}(\mu) \prod_{j=1}^{\ell(\mu)} \mu_j$, and $\chi_\mu(\nu)$ denotes the value of the irreducible character of S_d associated to $\nu \vdash d$ at the conjugacy class of S_d associated to $\mu \vdash d$.

Proposition 3.1.

$$
\begin{aligned}
&\sqrt{-1}^{\ell(\vec{\mu})}(-1)^{\sum_{i=1}^s f_i(|\mu^i|-1)}F^{\bullet\hat{Y},\hat{D}}_{\vec{d}',\vec{\mu}}(\lambda)\\
=&\sum_{|\nu^i|=|\mu^i|} G^{\bullet\hat{X}}_{\vec{d},\vec{\nu}}(\lambda;\mathsf{u},\mathsf{v})\prod_{i=1}^s z_{\nu^i}\Phi^{\bullet}_{\nu^i,\mu^i}(\sqrt{-1}\frac{\mathsf{f}_i}{\mathsf{w}_1^i})\\
=&\sum_{|\nu^i|=|\mu^i|} G^{\bullet\hat{X}}_{\vec{d},\vec{\nu}}(\lambda;\mathsf{u},\mathsf{v})\prod_{i=1}^s z_{\nu^i}\Phi^{\bullet}_{\nu^i,\mu^i}(\sqrt{-1}(f_i-\frac{\mathsf{w}_2^i}{\mathsf{w}_1^i})),
\end{aligned}
\tag{16}
$$

where $\vec{d}\colon E_c(\Gamma_X)\to\mathbb{Z}_{\geq 0}$ *is the restriction of* $\vec{d}'\colon E_c(\Gamma)\to\mathbb{Z}_{\geq 0}$.

Proof. We use the notation in [29, Section 7], and set $\nu^i=\nu^{\bar{e}_i}$. By [29, Proposition 7.10],

$$
\begin{aligned}
F^{\bullet\hat{Y},\hat{D}}_{\vec{d}',\vec{\mu}}(\lambda) \;=\;& \sum_{|\nu^{\bar{e}}|=\vec{d}'(\bar{e})}\prod_{\bar{e}\in E(\Gamma)}(-1)^{n^e\vec{d}'(\bar{e})}z_{\nu^{\bar{e}}}\prod_{v\in V_3(\Gamma)}\sqrt{-1}^{\ell(\vec{\nu}^v)}G^{\bullet}_{\vec{\nu}^v}(\lambda;\mathbf{w}_v)\\
&\cdot\prod_{i=1}^s(-1)^{|\mu^i|}(-\sqrt{-1})^{\ell(\nu^i)+\ell(\mu^i)}\Phi^{\bullet}_{\nu^i,\mu^i}(\sqrt{-1}\frac{\mathsf{f}_i}{\mathsf{w}_1^i}\lambda)\\
=\;&\sum_{|\nu^i|=|\mu^i|}\left(\sum_{|\nu^{\bar{e}}|=\vec{d}(\bar{e})}\prod_{\bar{e}\in E(\Gamma_X)}(-1)^{n^e\vec{d}(\bar{e})}z_{\nu^{\bar{e}}}\prod_{v\in V_3(\Gamma)}\sqrt{-1}^{\ell(\vec{\nu}^v)}G^{\bullet}_{\vec{\nu}^v}(\lambda;\mathbf{w}_v)\right)\\
&\prod_{i=1}^s(-1)^{(f_i-1)|\mu^i|}(-\sqrt{-1})^{\ell(\nu^i)+\ell(\mu^i)}z_{\nu^i}\Phi^{\bullet}_{\nu^i,\mu^i}(\sqrt{-1}\frac{\mathsf{f}_i}{\mathsf{w}_1^i}\lambda)
\end{aligned}
$$

where

$$
\sum_{|\nu^{\bar{e}}|=\vec{d}(\bar{e})}\prod_{\bar{e}\in E(\Gamma_X)}(-1)^{n^e}d^{\bar{e}}z_{\nu^{\bar{e}}}\prod_{v\in V_3(\Gamma)}\sqrt{-1}^{\ell(\vec{\nu}^v)}G^{\bullet}_{\vec{\nu}^v}(\lambda;\mathbf{w}_v)=\sqrt{-1}^{\ell(\vec{\nu})}G^{\bullet\hat{X}}_{\vec{d},\vec{\nu}}(\lambda;\mathsf{u},\mathsf{v})
$$

by localization computations similar to those in [29]. Therefore,

$$
F^{\bullet\hat{Y},\hat{D}}_{\vec{d}',\vec{\mu}}(\lambda)=(-\sqrt{-1})^{\ell(\vec{\mu})}(-1)^{\sum_{i=1}^s(f_i-1)|\mu^i|}\sum_{|\nu^i|=|\mu^i|}G^{\bullet\hat{X}}_{\vec{d},\vec{\nu}}(\lambda;\mathsf{u},\mathsf{v})z_{\nu^i}\Phi^{\bullet}_{\nu^i,\mu^i}(\sqrt{-1}\frac{\mathsf{f}_i}{\mathsf{w}_1^i}\lambda)
$$

□

3.4. **Open GW invariants of (X,L): multiple outer branes.** The open GW invariants of X relative to the framed outer branes $(L_1,f_1),\dots,(L_k,f_k)$ are defined to be

$$
N^{X,L}_{g,\beta',\mu^1,\dots,\mu^s}(f_1,\dots,f_s)=(-1)^{\sum_{i=1}^s(|\mu^i|-\ell(\mu^i))}\sum_{\pi(\vec{d}')=\beta'}F^{\hat{Y},\hat{D}}_{g,\vec{d}',\vec{\mu}}\in\mathbb{Q},
\tag{17}
$$

where $\vec{\mu}(v_i')=\mu^i$ for $i=1,\dots,s$, and the sign $(-1)^{\sum_{i=1}^s(|\mu^i|-\ell(\mu^i))}$ is included for convenience of comparison with the predictions from mirror symmetry.

Given $\beta\in H_2(X;\mathbb{Z})$ and $\vec{\mu}=(\mu^1,\dots,\mu^s)$, define

$$
\begin{aligned}
G^X_{g,\beta,\vec{\mu}}(\mathsf{u},\mathsf{v})=&\frac{(-\sqrt{-1})^{\ell(\vec{\mu})}}{|\mathrm{Aut}(\vec{\mu})|}\prod_{i=1}^s\prod_{j=1}^{\ell(\mu^i)}\frac{\prod_{m=1}^{\mu_j^i-1}(\mu_j^i\mathsf{w}_2^i+m\mathsf{w}_1^i)}{\mu_j^i\cdot\mu_j^i!(\mathsf{w}_1^i)^{\mu_j^i-1}}\\
&\cdot\int_{[\overline{\mathcal{M}}_{g,\ell(\vec{\mu})}(X,\beta)^{\mathbb{T}'}]^{\mathrm{vir}}}\frac{1}{e_{\mathbb{T}'}(N^{\mathrm{vir}})}\prod_{i=1}^s\prod_{j=1}^{\ell(\mu^i)}\frac{(\mathrm{ev}_j^i)^*\phi_i}{\frac{\mathsf{w}_1^i}{\mu_j^i}(\frac{\mathsf{w}_1^i}{\mu_j^i}-\psi_j^i)}
\end{aligned}
\tag{18}
$$

Then

$$G^X_{g,\beta,\vec{\mu}} = \sum_{\pi(\vec{d})=\beta} G^{\hat{X}}_{g,\vec{d},\vec{\mu}}$$

In particular, when $X=\mathbb{C}^3$, $s=3$, we have

$$\mathsf{w}_1^1=\mathsf{w}_2^3=\mathsf{u},\quad \mathsf{w}_2^1=\mathsf{w}_1^2=\mathsf{v},\quad \mathsf{w}_2^2=\mathsf{w}_1^3=-\mathsf{u}-\mathsf{v},$$

$$\overline{\mathcal{M}}_{g,\ell(\vec{\mu})}(\mathbb{C}^3,0)^{\mathbb{T}'}=\overline{\mathcal{M}}_{g,\ell(\vec{\mu})},$$

$$\frac{1}{e_{\mathbb{T}'}(N^{\mathrm{vir}})}=\prod_{i=1}^3\frac{\Lambda_g^\vee(\mathsf{w}_i)}{\mathsf{w}_i}\in H^*(\overline{\mathcal{M}}_{g,\ell(\vec{\mu})};\mathbb{Q})\otimes\mathbb{Q}(\mathsf{u},\mathsf{v}),$$

where $\Lambda_g^\vee(u)=u^g-\lambda_1u^{g-1}+\cdots+(-1)^g\lambda_g$. Define $\mathsf{w}_1=\mathsf{w}_4=\mathsf{u}$, $\mathsf{w}_2=\mathsf{v}$, and $\mathsf{w}_3=-\mathsf{u}-\mathsf{v}$. Then

$$\begin{aligned}G^{\mathbb{C}^3}_{g,0,\vec{\mu}}(\mathsf{u},\mathsf{v}) &= \frac{(-\sqrt{-1})^{\ell(\vec{\mu})}}{|\mathrm{Aut}(\vec{\mu})|}\prod_{i=1}^3\prod_{j=1}^{\ell(\mu^i)}\frac{\prod_{m=1}^{\mu^i_j-1}(\mu^i_j\mathsf{w}_{i+1}+m\mathsf{w}_i)}{\mu^i_j\cdot\mu^i_j!\mathsf{w}_i^{\mu^i_j-1}}\\ &\quad\cdot\int_{\overline{\mathcal{M}}_{g,\ell(\vec{\mu})}}\prod_{i=1}^3\frac{\Lambda_g^\vee(\mathsf{w}_i)\mathsf{w}_i^{\ell(\vec{\mu})-1}}{\prod_{m=1}^{\mu^i_j}\frac{\mathsf{w}_i}{\mu^i_j}(\frac{\mathsf{w}_i}{\mu^i_j}-\psi^i_j)}\end{aligned}$$

which is exactly the three-partition Hodge integral defined in [29, Section 2.3].

Given $\beta'\in H_2(X,L;\mathbb{Z})$ and an s-uple of partitions $\vec{\mu}=(\mu^1,\dots,\mu^s)$, define

$$F^{\bullet X,L}_{\beta',\vec{\mu}}(\lambda;f_1,\dots,f_s)=(-1)^{\sum_{i=1}^s(|\mu^i|-\ell(\mu^i))}\sum_{\pi(\vec{d}')=\beta'}F^{\bullet\hat{Y},\hat{D}}_{\vec{d}',\vec{\mu}}(\lambda).$$

The following proposition follows from Proposition 3.1 and definitions.

Proposition 3.2 (multiple framed outer branes)**.** *Let $\beta\in H_2(X;\mathbb{Z})$ be an effective curve class, and let $\beta'=\beta+\sum_{i=1}^s|\mu^i|b_i\in H_2(X,L;\mathbb{Z})$. Then*

$$\begin{aligned}&(-\sqrt{-1})^{\ell(\vec{\mu})}(-1)^{\sum_{i=1}^s f_i|\mu^i|}F^{\bullet X,L}_{\beta',\vec{\mu}}(\lambda;f_1,\dots,f_s)\\ &=\sum_{|\nu^i|=|\mu^i|}G^{\bullet X}_{\beta,\vec{\nu}}(\lambda;\mathsf{u},\mathsf{v})\prod_{i=1}^s z_{\nu^i}\Phi^\bullet_{\nu^i,\mu^i}(\sqrt{-1}\frac{\mathsf{f}_i}{\mathsf{w}_1^i})\\ &=\sum_{|\nu^i|=|\nu^i|}G^{\bullet X}_{\beta,\vec{\nu}}(\lambda;\mathsf{u},\mathsf{v})\prod_{i=1}^s z_{\nu^i}\Phi^\bullet_{\nu^i,\mu^i}(\sqrt{-1}(f_i-\frac{\mathsf{w}_2^i}{\mathsf{w}_1^i})).\end{aligned}\tag{19}$$

In particular, when $s=1$, $\mu^1=\mu$, $f_1=f$, $b_1=b$, we have

$$F^{\bullet X,L}_{\beta+|\mu|b,\mu}(\lambda;f)=\sqrt{-1}^{\ell(\mu)}(-1)^{f|\mu|}\sum_{|\nu|=|\mu|}G^{\bullet X}_{\beta,\nu}(\lambda;\mathsf{u},\mathsf{v})z_\nu\Phi^\bullet_{\nu,\mu}(\sqrt{-1}(f-\frac{\mathsf{v}_L}{\mathsf{u}_L})).$$

Let $\mathbb{T}_{L,f}\cong\mathbb{C}^*$ be the subtorus of $\mathbb{T}'$ defined in Section 2.5. It is the kernel of the character $\mathsf{v}_L-f\mathsf{u}_L\in\mathrm{Hom}(\mathbb{T}',\mathbb{C}^*)$. The inclusion $\mathbb{T}_{L,f}\hookrightarrow\mathbb{T}'$ induces a surjective ring homomorphism

$$H^*(B\mathbb{T}';\mathbb{Q})=\mathbb{Q}[\mathsf{u},\mathsf{v}]=\mathbb{Q}[\mathsf{u}_L,\mathsf{v}_L]\to H^*(\mathbb{T}_{L,f};\mathbb{Z})=\mathbb{Q}[\mathsf{u}_L]$$

given by $\mathsf{u}_L\mapsto\mathsf{u}_L$ and $\mathsf{v}_L\mapsto f\mathsf{u}_L$.

Recall that

$$\Phi^\bullet_{\nu,\mu}(0)=\frac{\delta_{\nu,\mu}}{z_\nu},$$

so

$$F^{\bullet X,L}_{\beta+|\mu|b,\mu}(\lambda;f)=\sqrt{-1}^{\ell(\mu)}(-1)^{f|\mu|}\sum_{|\nu|=|\mu|}G^{\bullet X}_{\beta,\nu}(\lambda;\mathsf{u},\mathsf{v})\Big|_{\mathsf{v}_L=f\mathsf{u}_L}.$$

Therefore,

$$N^{X,L}_{g,\beta+|\mu|b,\mu}(f)=\sqrt{-1}^{\ell(\mu)}(-1)^{f|\mu|}G^{X}_{\beta,\mu}(\lambda;\mathsf{u},\mathsf{v})|_{\mathsf{v}_L=f\mathsf{u}_L},$$

which is equivalent to the following corollary.

Corollary 3.3 (single framed outer brane)**.** *Let (L,f) be a framed outer brane in X, let z be the unique $\mathbb{T}'$-fixed point in $V(\tau_L)$, and let $\phi\in H^6_{\mathbb{T}'}(X;\mathbb{Z})$ be the $\mathbb{T}'$-equivariant Poincaré dual of z. For any effective curve class $\beta\in H_2(X;\mathbb{Z})$ and any partition $\mu=(\mu_1,\dots,\mu_h)$, we have*

$$\begin{aligned}N^{X,L}_{g,\beta+|\mu|b,\mu}(f)=&\frac{1}{|\mathrm{Aut}(\mu)|}\prod_{j=1}^{h}(-1)^{f\mu_j}\frac{\prod_{m=1}^{\mu_j-1}(f\mu_j+m)}{\mu_j\cdot\mu_j!}\\&\cdot\left(\int_{[\overline{\mathcal{M}}_{g,h}(X,\beta)^{\mathbb{T}'}]^{\mathrm{vir}}}\frac{1}{e_{\mathbb{T}'}(N^{\mathrm{vir}})}\prod_{j=1}^{h}\frac{\mathrm{ev}_j^*\phi}{\frac{\mathsf{u}_L}{\mu_j}(\frac{\mathsf{u}_L}{\mu_j}-\psi_j^i)}\right)\Big|_{\mathsf{v}_L=f\mathsf{u}_L}\end{aligned}\tag{20}$$

3.5. **Open GW invariants of (X,L): single outer or inner brane.** Let (L,f) be a framed (inner or outer) Aganagic-Vafa A-brane. Then $L=L_1$. Let

$$b=b_1\in H_2(X,L;\mathbb{Z}),\quad \gamma=\gamma_1\in H_1(L;\mathbb{Z}),\quad x^+=x_1,$$

where b_1,γ_1,x_1 are defined as in Section 2.6 Let $\phi_L^+\in H^6_{\mathbb{T}'}(X;\mathbb{Z})$ be the $\mathbb{T}'$-equivariant Poincaré dual of the $\mathbb{T}'$ fixed points $z^+\in V(\tau_L)$. When L is an inner brane, let $x^-=x_1^-$ be the other $\mathbb{T}'$ fixed point in $V(\tau_L)$, and let ϕ_L^- be the $\mathbb{T}'$-equivariant Poincaré dual of x^-.

Given $\beta'\in H_2(X,L;\mathbb{Z})$, and $\vec{w}=(w_1,\dots,w_h)$, where w_j are nonzero integers, we define

$$\overline{\mathcal{M}}_{g,\beta',\vec{w}}:=\overline{\mathcal{M}}_{g;h}(X,L\mid\beta';w_1\gamma,\dots,w_h\gamma)$$

where the right hand side is the moduli space of stable maps $u:(\Sigma,\partial\Sigma=\cup_{j=1}^h R_j)\to(X,L)$, where Σ is a prestable Riemann surface of type $(g;h)$, $R_j\cong S^1$ are connected components of the boundary $\partial\Sigma$ of Σ, $u_*[\Sigma]=\beta'$, $u_*[R_j]=w_j\gamma$. (See [20, Section 4] for the detailed definitions.) Then $\mathbb{T}'_{\mathbb{R}}$ acts on $\overline{\mathcal{M}}_{g,\beta',\vec{w}}$. The integral

$$\left(\int_{[\overline{\mathcal{M}}^{\mathbb{T}'_{\mathbb{R}}}_{g,\beta',\vec{w}}]^{\mathrm{vir}}}\frac{1}{e_{\mathbb{T}'_{\mathbb{R}}}(N^{\mathrm{vir}})}\right)\Big|_{\mathsf{v}_L=f\mathsf{u}_L}.$$

is a rational number depending on $f\in\mathbb{Z}$. It is defined up to a sign depending on choice of orientation on the virtual tangent bundle of $\overline{\mathcal{M}}_{g,\beta',\vec{w}}$. Define

$$\mathrm{Aut}(\vec{w})=\{\sigma\in S_h\mid(w_{\sigma(1)},\dots,w_{\sigma(h)})=(w_1,\dots,w_h)\}.$$

Proposition 3.4. *Suppose that $\beta\in H_2(X;\mathbb{Z})$, and either*

- *L is an outer brane, $w_1,\dots,w_h$ are positive integers, or*
- *L is an inner brane, $w_1,\dots,w_h$ are nonzero integers.*

When L is an inner brane, let $\alpha=[V(\tau_L)]\in H_2(X;\mathbb{Z})$. $\{1,\dots,h\}$ is a disjoint union of J_+ and J_-, where $J_\pm=\{j\in\{1,\dots,h\}\mid\pm w_j>0\}$. (So J_- is empty when L is an outer brane.) Define $\beta'\in H_2(X,L;\mathbb{Z})$ by

$$\beta'=\begin{cases}\beta+\left(\sum_{j=1}^h w_j\right)b & \text{if } L \text{ is an outer brane,}\\ \beta+\left(\sum_{j\in J_+}w_j\right)b+\left(\sum_{j\in J_-}(-w_j)\right)(\alpha-b) & \text{if } L \text{ is an inner brane.}\end{cases}$$

Then

$$\frac{1}{|\mathrm{Aut}(\vec{w})|}\int_{[\overline{\mathcal{M}}^{\mathbb{T}'_{\mathbb{R}}}_{g,\beta',\vec{w}}]^{\mathrm{vir}}}\frac{1}{e_{\mathbb{T}'_{\mathbb{R}}}(N^{\mathrm{vir}})}\Big|_{\mathsf{v}_L=f\mathsf{u}_L}=\pm N_{g,\beta'\vec{w}}(f)$$

where

$$\begin{aligned}N_{g,\beta',\vec{w}}(f)=&\frac{1}{|\mathrm{Aut}(\vec{w})|}\prod_{j\in J_+}(-1)^{fw_j}\frac{\prod_{m=1}^{w_j-1}(fw_j+m)}{w_j\cdot w_j!}\\ &\cdot\prod_{j\in J_-}(-1)^{(f+n)w_j}\frac{\prod_{m=1}^{-w_j-1}((f+n)(-w_j)+m)}{(-w_j)\cdot(-w_j)!}\\ &\cdot\left(\int_{[\overline{\mathcal{M}}_{g,h}(X,\beta)^{\mathbb{T}'}]^{\mathrm{vir}}}\frac{1}{e_{\mathbb{T}'}(N^{\mathrm{vir}})}\cdot\frac{\prod_{j\in J_+}\mathrm{ev}_j^*\phi_L^+\prod_{j\in J_-}\mathrm{ev}_j^*\phi_L^-}{\prod_{j=1}^h[\frac{\mathsf{u}}{w_j}(\frac{\mathsf{u}}{w_j}-\psi_j)]}\right)\Big|_{\mathsf{v}_L=f\mathsf{u}_L}\end{aligned}\tag{21}$$

Remark 3.5. The above formula (21) agrees with (20) when L is an outer brane. We use (21) to extend the definition of open GW invariants when (L,f) is an inner brane.

Proof of Proposition 3.4. When all the w_j's are positive, the computation for an inner brane is the same as that for an outer brane. From now on, we assume that L is an inner brane. $V(\tau_L)$ is the union of two disks D_+ and D_- which contain the torus fixed points x^+ and x^-, respectively. $D_+\cap D_-=L\cap V(\tau_L)$.



FIGURE 4. weights of the action by $T_{L,f}$

Suppose that $u:(C,q_1,\dots,q_h)\to X$ is a stable map which represents a $\mathbb{T}'$ fixed point in $\overline{\mathcal{M}}_{g,h}(X,\beta)$, and that $f(q_j)=x^{\pm}$ for $j\in J_{\pm}$. Let Σ be the prestable bordered Riemann surface

$$\Sigma=C\cup D_1\cup\cdots\cup D_h,$$

where D_j intersects C at a node q_j. If $j\in J_{\pm}$ then we define $u_j:D_j\to D_{\pm}$ by $z\mapsto z^{\pm w_j}$. Then we obtain a stable map $\hat{u}:(\Sigma,\partial\Sigma)\to(X,L)$ which represents a $\mathbb{T}'_{\mathbb{R}}$-fixed point in $\overline{\mathcal{M}}_{g,\beta',\vec{w}}$. Conversely, all $\mathbb{T}'_{\mathbb{R}}$ fixed points in $\overline{\mathcal{M}}_{g,\beta',\vec{w}}$ arise in this way. Let

$$\begin{aligned}&\hat{B}_1=\mathrm{Aut}(\Sigma),\ \hat{B}_2=\mathrm{Def}(\hat{u}),\ \hat{B}_4=\mathrm{Def}(\Sigma),\ \hat{B}_5=\mathrm{Obs}(\hat{u}),\\ &B_1=\mathrm{Aut}(C,q_1,\dots,q_h),\ B_2=\mathrm{Def}(u)\\ &B_4=\mathrm{Def}(C,q_1,\dots,q_h),\ B_5=\mathrm{Obs}(u).\end{aligned}$$

(See [20, Section 4] for the definitions of these real vector spaces.) Let $\hat{T}^1-\hat{T}^2$ be the virtual tangent space to $\overline{\mathcal{M}}_{g,\beta',\vec{w}}$ at the moduli point $[\hat{u}:(\Sigma,\partial\Sigma)\to(X,L)]$, and let T^1-T^2 be the virtual tangent space to $\overline{\mathcal{M}}_{g,h}(X,\beta)$ at the moduli point $[u:(C,q_1,\dots,q_h)\to X]$. Let $e=e_{\mathbb{T}'_{\mathbb{R}}}$ denote the $\mathbb{T}'_{\mathbb{R}}$-equivariant Euler class. Then

$$\frac{e(\hat{T}^{2,m})}{e(\hat{T}^{1,m})}=\frac{e(\hat{B}_1^m)e(\hat{B}_5^m)}{e(\hat{B}_4^m)e(\hat{B}_2^m)},\quad\frac{e(T^{2,m})}{e(T^{1,m})}=\frac{e(B_1^m)e(B_5^m)}{e(B_4^m)e(B_2^m)}$$

We have

$$\hat{B}_1^m = B_1^m, \quad \hat{B}_4^m = B_4^m \oplus \bigoplus_{j=1}^h T_{q_j} C \otimes T_{q_j} D_j.$$

Therefore

$$\frac{e(\hat{B}_1^m)}{e(\hat{B}_4^m)} = \frac{e(B_1^m)}{e(B_4^m)} \prod_{j\in J_+} (\frac{\mathsf{u}_L}{w_j} - \psi_j) \prod_{j\in J_-} (\frac{-\mathsf{u}_L}{-w_j} - \psi_j) = \frac{e(B_1^m)}{e(B_4^m)} \prod_{j=1}^h (\frac{\mathsf{u}_L}{w_j} - \psi_j) \tag{22}$$

We also have a long exact sequence

$$\begin{aligned} 0 \quad &\to \hat{B}_2 \to B_2 \oplus \textstyle\bigoplus_{j=1}^h H^0(D_j) \to \bigoplus_{j\in J_+} T_{x^+}X \oplus \bigoplus_{j\in J_-} T_{x^-}X \\ &\to \hat{B}_5 \to B_5 \oplus \textstyle\bigoplus_{j=1}^h H^1(D_i) \to 0. \end{aligned}$$

where

$$H^p(D_j) = H^p(D_j, \partial D_j, u_j^* TX, (u_j|_{\partial D_j})^* TL), \quad p = 0, 1.$$

We have

$$\begin{aligned} e_{\mathbb{T}'}(T_{x^+}X) &= \mathsf{u}_L \mathsf{v}_L (-\mathsf{u}_L - \mathsf{v}_L) = \mathrm{ev}_j^* \phi_L^+ \quad j \in J_+, \\ e_{\mathbb{T}'}(T_{x^-}X) &= (-\mathsf{u}_L)(-\mathsf{v}_L - n\mathsf{u}_L)(\mathsf{v}_L + (n+1)\mathsf{u}) = \mathrm{ev}_j^* \phi_L^- \quad j \in J_- \\ \frac{e(H^1(D_j)^m)}{e(H^0(D_j)^m)} &= \pm \frac{1}{\mathsf{u}_L} \prod_{m=1}^{w_j - 1} \frac{w_j \mathsf{v}_L + m\mathsf{u}_L}{m\mathsf{u}_L}, \quad j \in J_+ \\ \frac{e(H^1(D_j)^m)}{e(H^0(D_j)^m)} &= \pm \frac{1}{\mathsf{u}_L} \prod_{m=1}^{w_j - 1} \frac{w_j \mathsf{v}_L + (n+m)\mathsf{u}_L}{m\mathsf{u}_L}, \quad j \in J_- \end{aligned}$$

Therefore

$$\frac{e(\hat{B}_5^m)}{e(\hat{B}_2^m)} = \frac{e(B_5^m)}{e(B_2^m)} \cdot \prod_{j\in J_+} \frac{\mathrm{ev}_j^* \phi_+^L}{\mathsf{u}_L} \prod_{m=1}^{w_j-1} \frac{w_j \mathsf{v}_L + m\mathsf{u}_L}{m\mathsf{u}_L} \cdot \prod_{j\in J_-} \frac{\mathrm{ev}_j^* \phi_-^L}{-\mathsf{u}_L} \prod_{m=1}^{w_j-1} \frac{w_j \mathsf{v}_L + (n+m)\mathsf{u}_L}{m\mathsf{u}_L} \tag{23}$$

Finally,

$$\mathrm{Aut}(\hat{u}) = \mathrm{Aut}(u) \times \prod_{j=1}^h (\mathbb{Z}/w_j\mathbb{Z}). \tag{24}$$

Combining (22), (23), and (24), we get

$$\begin{aligned} N_{g,\beta',\vec{w}}(f) = \pm \frac{1}{|\mathrm{Aut}(\vec{w})|} \prod_{j\in J_+} \frac{\prod_{m=1}^{w_j-1}(fw_j + m)}{w_j \cdot w_j!} \prod_{j\in J_-} \frac{\prod_{m=1}^{-w_j-1}((f+n)(-w_j)+m)}{(-w_j)\cdot(-w_j)!} \\ \cdot \Big(\int_{[\overline{\mathcal{M}}_{g,h}(X,\beta)^{\mathbb{T}'}]^{\mathrm{vir}}} \frac{1}{e_{\mathbb{T}'}(N^{\mathrm{vir}})} \cdot \frac{\prod_{j\in J_+} \mathrm{ev}_j^* \phi_L^+ \prod_{j\in J_-} \mathrm{ev}_j^* \phi_L^-}{\prod_{j=1}^h [\frac{\mathsf{u}_L}{w_j}(\frac{\mathsf{u}_L}{w_j} - \psi_j)]} \Big) \Big|_{\mathsf{v}_L = f\mathsf{u}_L} \end{aligned}$$

□

3.6. **Generating functions for a fixed topological type of the domain.** Any $\beta' \in H_2(X, L; \mathbb{Z})$ is of the form $\beta' = \beta + wb$, where $\beta \in H_2(X; \mathbb{Z})$ and $w \in \mathbb{Z}$. As in Section 2, we fix a choice of $\mathbb{Z}$-basis $l^{(1)}, \ldots, l^{(k)}$ and its dual $p_1, \ldots, p_k$. Define

$$Q^{\beta'} = Q_0^w \prod_{a=1}^{k} Q_a^{d_a}, \quad d_a = \langle \beta, p_a \rangle.$$

We define generation functions

$$\begin{aligned} F_{g,\vec{w}}(Q; f) &= \sum_{\beta' \in H_2(X,L;\mathbb{Z})} N_{g,\beta',\vec{w}}(f) Q^{\beta'}, \\ F^{g,h}(Q, Y; f) &= \sum_{w_j \in \mathbb{Z} - \{0\}} F_{g,w_1,\ldots,w_h}(Q, f) \prod_{j=1}^{h} Y_j^{w_j}, \end{aligned}$$

where

$$Q = (Q_0, Q_1, \ldots, Q_k), \quad Y = (Y_1, \ldots, Y_h).$$

Note that when $h = 1$, $w = w_1 \in \mathbb{Z}$ is determined by $\partial \beta' = w\gamma$, so we may omit the variable Y_1. In particular, we define

$$F(Q; f) := F^{0,1}(Q, Y_1 = 1; f) = \sum_{w \in \mathbb{Z}, w \neq 0, \beta \in H_2(X;\mathbb{Z})} N_{0,\beta+wb,w} Q_0^w \prod_{a=1}^{k} Q_a^{d_a}, \tag{25}$$

where $d_a = \langle \beta, p_a \rangle$.

For later convenience, we introduce the following generating function for any nonzero integer w:

$$J_w^L(Q; f) = \begin{cases} 1 + \sum_{\beta > 0} Q^\beta \left(\int_{[\overline{\mathcal{M}}_{0,1}(X,\beta)^{\mathbb{T}'}]^{\mathrm{vir}}} \frac{1}{e_{\mathbb{T}'}(N^{\mathrm{vir}})} \frac{\mathrm{ev}_j^* \phi_L^+}{\frac{\mathfrak{u}_L}{w}(\frac{\mathfrak{u}_L}{w} - \psi_j)]} \right) \Big|_{\mathfrak{v}_L = f\mathfrak{u}_L} & w > 0 \\ 1 + \sum_{\beta > 0} Q^\beta \left(\int_{[\overline{\mathcal{M}}_{0,1}(X,\beta)^{\mathbb{T}'}]^{\mathrm{vir}}} \frac{1}{e_{\mathbb{T}'}(N^{\mathrm{vir}})} \frac{\mathrm{ev}_j^* \phi_L^-}{\frac{\mathfrak{u}_L}{w}(\frac{\mathfrak{u}_L}{w} - \psi_j)]} \right) \Big|_{\mathfrak{v}_L = f\mathfrak{u}_L} & w < 0 \end{cases} \tag{26}$$

where $\beta \in H_2(X; \mathbb{Z})$. We say $\beta \geq 0$ when β is effective, and $\beta > 0$ when it is also non-zero. The following is an immediate consequence of Proposition 3.4.

Corollary 3.6. (a) *If L is an outer brane then*

$$F(Q; f) = \sum_{w > 0} Q_0^w (-1)^{fw} \frac{\prod_{m=1}^{w-1}(fw + m)}{w \cdot w!} J_w^L(Q; f). \tag{27}$$

(b) *If is an inner brane then*

$$\begin{aligned} F(Q; f) = &\sum_{w > 0} Q_0^w (-1)^{fw} \frac{\prod_{m=1}^{w-1}(fw + m)}{w \cdot w!} J_w^L(Q; f) \\ &+ \sum_{w < 0} (\frac{Q_0}{Q^\alpha})^w (-1)^{(f+n)w} \frac{\prod_{m=1}^{-w-1}((f+n)(-w) + m)}{-w \cdot (-w)!} J_w^L(Q; f) \end{aligned} \tag{28}$$

4. The Mirror Conjecture on Holomorphic Disks

Aganagic-Vafa and Aganagic-Klemm-Vafa [3, 2] construct a B-model Landau-Ginzburg model together with the superpotential W as the mirror to the A-model on the Calabi-Yau toric threefold X. The superpotential W is conjectured, up to a mirror transform, to be equal to the single brane disk amplitude constructed in Section 3.6. As discussed in Section 3.5 we fix a framed Aganagic-Vafa A-brane (L, f), and let x^+ be the $\mathbb{T}$-fixed point in $V(\tau_L)$ (when L is an inner brane there is another $\mathbb{T}$-fixed point x^-). We fix indices i_1, i_2, i_3, and in case of inner brane L, also an index i_4 as in Section 2.5 (Figure 4).

4.1. **The Hori-Vafa mirror.** Applying $\mathrm{Hom}(-, \mathbb{C}^*)$ to (5), we obtain

$$1 \to \mathbb{T}^\vee \to \widetilde{\mathbb{T}}^\vee \xrightarrow{\phi} G^\vee \to 1, \tag{29}$$

where $\mathbb{T}^\vee \cong (\mathbb{C}^*)^n$, $\widetilde{\mathbb{T}}^\vee \cong (\mathbb{C}^*)^r$, $G^\vee \cong (\mathbb{C}^*)^k$. Applying $\mathrm{Hom}(-, \mathbb{C}^*)$ to (9), we obtain

$$1 \to (\mathbb{C}^*)^\vee \to \mathbb{T}^\vee \to (\mathbb{T}')^\vee \to 1, \tag{30}$$

where $(\mathbb{T}')^\vee \cong (\mathbb{C}^*)^{n-1}$.

Given $q = (q_1, \dots, q_k) \in G^\vee$, define

$$\mathbb{T}_q^\vee := \phi^{-1}(q) = \{\vec{x} = (x_1, \dots, x_{k+3}) \in \widetilde{\mathbb{T}}^\vee \mid \prod_{i=1}^{k+3} x_i^{l_i^{(a)}} = q_a,\ a = 1, \dots, k\} \cong (\mathbb{C}^*)^3.$$

Let $\mathbb{C}^*$ acts on $\widetilde{\mathbb{T}}^\vee$ by $\lambda \cdot (x_1, \dots, x_{k+3}) = (\lambda x_1, \dots, \lambda x_{k+3})$. Then the $\mathbb{C}^*$-action preserves $\mathbb{T}_q^\vee$. Let $\mathbb{T}_q'^\vee := \mathbb{T}_q^\vee / \mathbb{C}^* \cong (\mathbb{C}^*)^2$.

Let $H(\vec{x}, q)$ be the restriction of $\sum_{i=1}^{k+3} x_i$ to $\mathbb{T}_q^\vee$. There is a $\mathbb{C}^*$ action on the following Calabi-Yau 4-fold

$$\widetilde{X}_q^\vee = \{(w^+, w^-, \vec{x}) \in \mathbb{C}^2 \times \mathbb{T}_q^\vee \mid w^+ w^- = H(\vec{x}, q)\},$$

given by

$$\begin{aligned} \lambda \in \mathbb{C}^* : \qquad w^- &\mapsto \lambda w^-, \\ (x_1, \dots, x_{k+3}) &\mapsto (\lambda x_1, \dots, \lambda x_{k+3}). \end{aligned}$$

The mirror of X is the quotient noncompact Calabi-Yau 3-fold $\widetilde{X}_q^\vee / \mathbb{C}^*$, denoted by $X_q^\vee$. The curve in $X_q^\vee$ given by the equation $H(\vec{x}, q) = 0$, $w^+ = w^- = 0$ is the *mirror curve*.

For $a = 1, \dots, k$, we extend the charge vectors $l^{(a)} = (l_1^{(a)}, \dots, l_{k+3}^{(a)})$ to

$$\widetilde{l}^{(a)} = (l_1^{(a)}, \dots, l_{k+3}^{(a)}, 0, 0).$$

There is an additional charge vector $\widetilde{l}^{(0)} = (l_1^{(0)}, \dots, l_{k+5}^{(0)})$ for $i = 1, \dots, k+5$ with

$$\sum_{i=1}^{k+5} l_i^{(0)} = 0$$

that characterizes the B-brane. Define the index sets

$$I = \{1, \dots, r\}, \quad \widetilde{I} = \{1, \dots, r+2\}, \quad I_0 = \{i_1, i_2, i_3\}. \tag{31}$$

If the A-brane carries the framing f, we set the open charge vector $l^{(0)} = (l_1^{(0)}, \dots, l_{k+3}^{(0)})$ to be

$$l_{i_1}^{(0)} = 1, \quad l_{i_2}^{(0)} = f, \quad l_{i_3}^{(0)} = -f - 1, \quad l_i^{(0)} = 0 \quad \text{for } i \in I \backslash I_0.$$

We introduce an extra charge vector

$$\widetilde{l}^{(0)} = (l_1^{(0)}, \dots, l_r^{(0)}, 1, -1).$$

The mirror curve can be written down in the following coordinate patch

$$x_{i_1} = -\widetilde{x},\ x_{i_2} = \widetilde{y},\ x_{i_3} = 1.$$

Other x_i for $i \in I$ are determined from $\widetilde{x}, \widetilde{y}$ through the equation

$$\prod_{i=1}^{k+3} x_i^{l_i^{(a)}} = q_a, \quad a = 1, \ldots, k.$$

The mirror curve is an affine curve in $\mathbb{C}^* \times \mathbb{C}^*$ given by a single equation

$$H(\widetilde{x}, q) = -\widetilde{x} + \widetilde{y} + 1 + \sum_{i \in I \setminus I_0} x_i(\widetilde{x}, \widetilde{y}, q) = 0$$

in coordinates $\widetilde{x}, \widetilde{y}$.

4.2. **Aganagic-Vafa B-branes.** As shown in [3], the mirror B-brane of the A-brane defined in Section 2.4 above is 2-cycle in $X_q^\vee$, given by the equations

$$w^- = 0,\ H(\vec{x}, q) = 0, \tag{32}$$

where $\vec{x} = \vec{x}(w^+) \in \{\vec{x} | H(\vec{x}, q) = 0\}$ is a function of the coordinate w^+ on the brane. We fix $\vec{x}$ at $w^+ = \infty$ to be $\vec{x}(\infty) = \vec{x}^*$. Following [23, 24], the framed modulus x of the B-brane is determined by the value of $\vec{x}$ on $w^+ = 0$

$$x = q_0 = -\prod_{i=1}^{r} x_i(0)^{l_i^{(0)}} = \widetilde{x}\widetilde{y}^f.$$

The the B-brane modulus in framing 0 is given by $\widetilde{x}$. We denote the corresponding B-brane by $\mathcal{C}_x$, and the reference B-brane by $\mathcal{C}_{x^*}$, with $x^* = -\prod_{i=1}^{k+3} x_i^*(0)^{l_i^{(0)}}$. The reference brane $\mathcal{C}_{x^*}$ could be chosen as a holomorphic curve with constant $x(w^+) = x^*$. Other branes cannot be holomorphic, since x has different bounded values at $w^+ = 0$ and $w^+ = \infty$.

The mirror curve in coordinates $\widetilde{x}, \widetilde{y}$ can be further written as an equation in the framed coordinates x, y

$$M(x, y, q) = H(\vec{x}, q) = -xy^{-f} + y + 1 + \sum_{i \in I \setminus I_0} x_i(xy^{-f}, y, q) = 0 \tag{33}$$

by a change of coordinates

$$x = \widetilde{x}\widetilde{y}^f,\ y = \widetilde{y}.$$

4.3. **Picard-Fuchs equations and the mirror map.** Let $\Gamma(x)$ be a 3-chain in $X_q^\vee$ such that $\partial\Gamma(x) = \mathcal{C}_x - \mathcal{C}_{x^*}$, and $\gamma \in H^3(X_q^\vee; \mathbb{Z})$ is any 3-cycle. Mayr and Lerche-Mayr [34, 23] show that the periods

$$\int_{\Gamma(x)} \Omega$$

are eliminated by GKZ-type operators

$$\mathcal{D}_a = \prod_{l_i^{(a)} > 0} \prod_{j=0}^{l_i^{(a)} - 1} (\sum_b l_i^{(b)} \theta_b - j) - q_a \prod_{l_i^{(a)} < 0} \prod_{j=0}^{l_i^{(a)} - 1} (\sum_{b=1}^{k} l_i^{(b)} \theta_b - j),$$

where $a = 0, \ldots, k$, $i = 1, \ldots, r+2$, and $\theta_a = q^a \frac{d}{dq^a}$. The extended Picard-Fuchs equations are

$$\mathcal{D}_\alpha W(q_0, q_1, \ldots, q_k) = 0. \tag{34}$$

Let $q_a = e^{t_a}$, and the string flat coordinates $Q_a = e^{T_a}$. The open string variable is $Q_0 = e^{T_0}$. Recall that for $a = 1, \ldots, k$, $T_a = -r_a + \sqrt{-1}\theta_a$ are complexified Kähler parameters. The mirror map is

$$T_a = \Pi_a^1(q).$$

where $\Pi_a^1(q)$ is a solution to the Picard-Fuchs equations above, i.e. $\mathcal{D}_b \Pi_a^1(q) = 0$ for $b = 0, \ldots, k$. It has the leading term behavior

$$T_a = \log(q_a) + S_a(q), \tag{35}$$

where $S_a(q)$ is a power series in $q_1, \ldots, q_k$.

Recall that $d_a = \langle p_a, \beta \rangle$ for $\beta \in H_2(X;\mathbb{Z})$, and $\langle D_i^*, \beta \rangle = \sum_{a=1}^k d_a l_i^{(a)}$. Denote $q^\beta = \prod_{a=1}^k q_a^{d_a}$. For any $\beta \in \mathbb{L}_{\text{eff}}$ and $i_0 \in I$, if $\langle D_{i_0}^*, \beta \rangle < 0$ and $\langle D_{i_0}^*, \beta \rangle \geq 0$ for $i \in I \backslash \{i_0\}$, we define

$$E_{i_0}(\beta) = \frac{(-1)^{(-\langle D_{i_0}^*, \beta \rangle - 1)}(-\langle D_{i_0}^*, \beta \rangle - 1)!}{\prod_{i \in I \backslash \{i_0\}} \langle D_i^*, \beta \rangle !};$$

otherwise we define $E_{i_0}(\beta) = 0$. Recall that $\beta \geq 0$ when β is effective, and $\beta > 0$ when it is also non-zero. In particular, using the Frobenius method, as shown in [36, 23], the mirror correction is given by

$$S_a(q) = \sum_{i \in I} l_i^{(a)} A_i(q), \quad a = 0, \ldots, k, \tag{36}$$

where $A(q)$ is the following series of $q_1, \ldots, q_k$:

$$A_i(q) = \sum_{\beta > 0} E_i(\beta) q^\beta. \tag{37}$$

The above formula is a direct reformulation of the equations [23, (2.20,2.21,2.22)].

Remark 4.1. In [36, 23], Lerche-Mayr explicitly compute the open mirror correction $S_0(q)$. Let $I' \subset \{1, \ldots, r\}$ be the index set such that $A_i(q) = 0$ for $i \notin I'$. One chooses an index subset $K \subset \{1, \ldots, k\}$ such that $|K| = |I'|$ and S_a are linearly independent. Denote the matrix $L = (l_i^{(a)})_{i \in I'}^{a \in K}$. Lerche-Mayr in [23] gives the open mirror correction as $S_0(q) = \sum_{i \in I'} l_i^{(0)} (\sum_{a \in K} (L^{-1})_a^i S_a(q))$. It turns out that

$$\begin{aligned} S_0(q) &= \sum_{i \in I'} l_i^{(0)} (\sum_{a \in K} (L^{-1})_a^i S_a(q)) \\ &= \sum_{i \in I'} l_i^{(0)} (\sum_{a \in K} (L^{-1})_a^i \sum_{j \in I'} l_j^{(a)} A_j(q)) \\ &= \sum_{i \in I'} l_i^{(0)} A_i(q). \end{aligned}$$

It agrees with (36).

4.4. **Superpotential and the mirror prediction of disk invariants.** In [3], the superpotential $W(\mathcal{C}_x)$ associated to the B-brane $\mathcal{C}_x$ is the integral over the chain

$$W = \int_{\Gamma(x)} \Omega,$$

where the boundary of the chain $\partial\Gamma(x) = \mathcal{C}_x - \mathcal{C}_{x^*}$. Let $\widetilde{\mathbb{L}} = \mathbb{Z}\times\mathbb{L} \cong H_2(X, L; \mathbb{Z})$. For any $\beta' = (w, \beta) \in \widetilde{\mathbb{L}}$, denote $q^{\beta'} = q_0^w q^\beta$. We define the extended pairing

$$\langle D_i^*, \beta'\rangle = w\widetilde{l}_i^{(0)} + \langle D_i^*, \beta\rangle = \sum_{a=0}^{k} d_a \widetilde{l}_i^{(a)}.$$

Define the cone $\widetilde{\mathbb{L}}_{\text{eff}}(L)$ in $\mathbb{Z}\times\mathbb{L}_{\text{eff}} \subset \widetilde{\mathbb{L}}$

$$\widetilde{\mathbb{L}}_{\text{eff}}(L) = \{\beta' = (w, \beta) \in \widetilde{\mathbb{L}} | w \neq 0, \langle D_i^*, \beta'\rangle \geq 0, i \in I\backslash\{i_2, i_3\}\}.$$

Given the charge vectors $\widetilde{l}^{(a)}$, $a = 0, \ldots, k$, Lerche and Mayr [36, 23] show that W is a double logarithm solution of the Picard-Fuchs equations. Precisely W consists of a double logarithm part and a power series part

$$W = \sum_{i,j=0}^{k} c_{ij} \log q_i \log q_j + W_0,$$

where W_0 is explicitly in the following form

$$W_0 = \sum_{\beta'\in\mathbb{L}_{\text{eff}}(L)} C(\beta') q^{\beta'}. \tag{38}$$

The coefficient $C(\beta')$ is obtained by applying the Frobenius method

$$\begin{aligned} C(\beta') &= \begin{cases} \dfrac{\prod_{i=i_3, r+3}(-1)^{-\langle D_i^*, \beta'\rangle - 1}(-\langle D_i^*, \beta'\rangle - 1)!}{\prod_{i\in\widetilde{I}\backslash\{i_3, r+3\}}\langle D_i^*, \beta'\rangle!}, & \langle D_{i_3}^*, \beta'\rangle < 0 \\ \dfrac{\prod_{i=i_2, r+3}(-1)^{-\langle D_i^*, \beta'\rangle - 1}(-\langle D_i^*, \beta'\rangle - 1)!}{\prod_{i\in\widetilde{I}\backslash\{i_2, r+3\}}\langle D_i^*, \beta'\rangle!}, & \langle D_{i_2}^*, \beta'\rangle < 0 \end{cases} \\ &= \frac{(-1)^{\langle D_{i_3}^* + D_{r+3}, \beta'\rangle}}{\prod_{i\in\widetilde{I}\backslash\{i_2, i_3, r+3\}}\langle D_i^*, \beta'\rangle!} \frac{\prod_{m=-\infty}^{-\langle D_{i_3}^*, \beta'\rangle - 1} m}{\prod_{m=-\infty}^{\langle D_{i_2}^*, \beta'\rangle} m}. \end{aligned}$$

In [3] and [2], the superpotential W_0 is explicitly computed by solving the mirror curve $M(x, y, q) = 0$. Writing $y = y(x, q)$, and in [3] it is shown that the superpotential is

$$x\frac{\partial W_0}{\partial x} = -\log y(x, q) \quad \text{up to a function in } q.$$

Remark 4.2. The above formula differs by a sign with Equation 4.5 of [3]. We choose this sign convention in order to match the sign convention on Gromov-Witten theory. One may simply apply a change of coordinate $y \mapsto 1/y$ to get rid of this minus sign. We only consider invariants of non-zero winding numbers. Bouchard and Sułkowski discuss the constant contributions of the higher genus superpotential associated to a mirror curve [6].

Conjecture 4.3. *After a change of variables by the mirror transform* (35)*,*

$$F(Q; f) = W_0(q; f).$$

We prove this conjecture in the next section.

5. Proof of Conjecture 4.3

5.1. **Equivariant mirror theorem.** A toric manifold X is semi-Fano if $c_1(T_X) \geq 0$. In this subsection, we state an equivariant mirror theorem for a semi-projective semi-Fano toric manifold X (Theorem 5.1). When X is projective, or is the total space of a direct sum of negative line bundles over a projective toric manifold Z, Theorem 5.1 follows from the results in [30, 13] (when X or Z is a projective space) and in [31, 14] (in the general case). When X is a general noncompact, semi-projective, semi-Fano toric manifold, Theorem 5.1 follows from the results in [9].

Let X be a semi-projective toric manifold. Let $D_1, \dots, D_r$ be the $\mathbb{T}$-divisor of X. Let

$$D_i^* = c_1(\mathcal{O}_X(D_i)) \in H^2(X;\mathbb{Z}), \quad (D_i^*)_{\mathbb{T}} = (c_1)_{\mathbb{T}}(\mathcal{O}_X(D_i)) \in H^*_{\mathbb{T}}(X;\mathbb{Z}).$$

Define an $H^*_{\mathbb{T}}(X;\mathbb{Z})$-valued function $I(t,z^{-1})$:

$$I(t,z^{-1}) = e^{(t_0'+\sum_{a=1}^k p_a t_a)/z}\Big(1+\sum_{\beta>0} q^\beta \prod_{i=1}^r \frac{\prod_{m=-\infty}^0((D_i^*)_{\mathbb{T}}+mz)}{\prod_{m=-\infty}^{\langle \beta, D_i^*\rangle}((D_i^*)_{\mathbb{T}}+mz)}\Big).$$

Note that the parameter t_0' is different from the parameter $t_0 = \log q_0$ introduced in Section 4.3.

Define another $H^*_{\mathbb{T}}(X;\mathbb{Z})$-valued function $J(T,z^{-1})$ as follows:

$$\begin{aligned} & \langle \gamma, J\rangle \\ = & \int_X e^{(T_0'+\sum_{a=1}^k p_a T_a)/z}\gamma + \sum_{\beta>0} Q^\beta \int_{[\overline{\mathcal{M}}_{0,2}(X,d)]^{\mathrm{vir}}} \frac{\mathrm{ev}_1^*(e^{(T_0'+p_1T_1+\cdots+p_kT_k)/z}\gamma)\mathrm{ev}_2^*(1)}{z-\psi_1} \\ = & \int_X e^{(T_0'+\sum_{a=1}^k p_a T_a)/z}\gamma + \sum_{\beta>0} Q^\beta \int_{[\overline{\mathcal{M}}_{0,1}(X,d)]^{\mathrm{vir}}} \frac{\mathrm{ev}_1^*(e^{(T_0'+\sum_{a=1}^k p_a T_a)/z}\gamma)}{z(z-\psi_1)}; \end{aligned}$$

or equivalently,

$$J = e^{(T_0'+\sum_{a=1}^k T_a p_a)/z}\Big(1+\sum_{\beta>0} Q^\beta (\mathrm{ev}_1^d)_* \frac{1}{z(z-\psi_1)}\Big).$$

Assume that X is semi-Fano. Then the expansion of the I function in z^{-1} is of the following form

$$I(t,z^{-1}) = 1 + z^{-1}(t_0' + f_0(q) + \sum_{i=1}^r \lambda_i g_i(q) + \sum_{a=1}^k p_a(t_a + f_a(q))) + O(z^{-2}),$$

where $\lambda_1, \dots, \lambda_r \in H^2(B\mathbb{T})$ are the universal first Chern classes. The expansion of the J function in z^{-1} is

$$J(T,z^{-1}) = 1 + z^{-1}(T_0' + \sum_{i=1}^k p_a T_a) + O(z^{-2}).$$

The mirror map is given by

$$T_0' = t_0' + f_0(q) + \sum_{i=1}^r \lambda_i g_i(q), \quad T_a = t_a + f_a(q), \quad a = 1, \dots, k.$$

In particular,

$$Q_a = q_a \exp(f_a(q)), \quad a = 1, \dots, k.$$

Theorem 5.1 below is a special case of the results in [9]. When X is the total space of a direct sum of negative line bundles over a projective toric manifold (such as the six toric Calabi-Yau 3-folds considered in Section 6.1–6.6), Theorem 5.1 is a special case of the results in [31, 14]. (It is stated as Corollary 4.3 in [14].)

Theorem 5.1 (Equivariant mirror theorem)**.** *With the above identification*

$$T_0' = t_0' + f_0(q) + \sum_{i=1}^{r} \lambda_i g_i(q), \quad T_a = t_a + f_a(q), \quad a = 1, \dots, k,$$

the I and J functions are equal

$$I(t, z^{-1}) = J(T, z^{-1}).$$

5.2. **Open and closed mirror maps.** We now restrict to the subtorus $\mathbb{T}_{L,f}$. Let $\iota_+ : x^+ \to X$ be the inclusion map, and let $\phi^+ \in H^6_{\mathbb{T}_{L,f}}(X;\mathbb{Z})$ be the $\mathbb{T}_{L,f}$-equivariant Poincaré dual of x^+. Recall that $J^L_w(Q;f)$ is defined by (26) and $A_i(q)$ is defined by in (37).

Proposition 5.2. *Let w be a positive integer. Then*

$$J^L_w(Q;f) = (\frac{q_0}{Q_0})^w \left(1 + \sum_{\beta>0} q^\beta \prod_{i=1}^{r} \frac{\prod_{m=-\infty}^{0}(w l_i^{(0)} + m)}{\prod_{m=-\infty}^{\langle \beta, D_i^* \rangle}(w l_i^{(0)} + m)}\right).$$

where $Q = (Q_0, Q_1, \dots, Q_k)$ and $q = (q_0, q_1, \dots, q_k)$ are related by the following open and closed mirror maps

$$Q_a = q_a \exp(\sum_{i \in I} l_i^{(a)} A_i(q)), \quad a = 0, 1, \dots, k. \tag{39}$$

Proof. In this proof, $\mathsf{u} = \mathsf{u}_L$. The parameters t_0' and t_0 are different.

$$\iota_+^* J(T, z^{-1}) = e^{(T_0' + \sum_{a=1}^k T_a \iota_+^* p_a)/z} \Big(1 + \sum_{\beta>0} Q^\beta \int_{[\overline{\mathcal{M}}_{0,1}(X,\beta)]^{\mathrm{vir}}} \frac{\mathrm{ev}_1^* \phi^+}{z(z - \psi_1)}\Big).$$

$$\begin{aligned}
\iota_+^* J(T, \frac{w}{\mathsf{u}}) &= e^{(T_0' + \sum_{a=1}^k T_a \iota_+^* p_a)\frac{w}{\mathsf{u}}} \Big(1 + \sum_{\beta>0} Q^\beta \int_{[\overline{\mathcal{M}}_{0,1}(X,\beta)]^{\mathrm{vir}}} \frac{\mathrm{ev}_1^* \phi^+}{\frac{\mathsf{u}}{w}(\frac{\mathsf{u}}{w} - \psi_1)}\Big) \\
&= e^{(T_0' + \sum_{a=1}^k T_a \iota_+^* p_a)\frac{w}{\mathsf{u}}} J^L_w(Q;f) \\
\iota_+^* I(t, \frac{w}{\mathsf{u}}) &= e^{(t_0' + \sum_{a=1}^k t_a \iota_+^* p_a)\frac{w}{\mathsf{u}}} \Big(1 + \sum_{\beta>0} q^\beta \prod_{i=1}^{r} \frac{\prod_{m=-\infty}^{0}(l_i^{(0)}\mathsf{u} + m\frac{\mathsf{u}}{w})}{\prod_{m=-\infty}^{\langle \beta, D_i^* \rangle}(l_i^{(0)}\mathsf{u} + m\frac{\mathsf{u}}{w})}\Big) \\
&= e^{(t_0' + \sum_{a=1}^k t_a \iota_+^* p_a)\frac{w}{\mathsf{u}}} \Big(1 + \sum_{\beta>0} q^\beta \prod_{i=1}^{r} \frac{\prod_{m=-\infty}^{0}(w l_i^{(0)} + m)}{\prod_{m=-\infty}^{\langle \beta, D_i^* \rangle}(w l_i^{(0)} + m)}\Big)
\end{aligned}$$

where we used $\sum_{i=1}^r D_i^* = 0$. We conclude that

$$J^L_w(Q;f) = (\frac{q_0}{Q_0})^w \left(1 + \sum_{\beta>0} q^\beta \prod_{i=1}^{r} \frac{\prod_{m=-\infty}^{0}(w l_i^{(0)} + m)}{\prod_{m=-\infty}^{\langle \beta, D_i^* \rangle}(w l_i^{(0)} + m)}\right).$$

where Q_0 and q_0 are related by the following open mirror map

$$Q_0 = q_0 \exp(\frac{T_0' - t_0'}{\mathsf{u}} + \sum_{a=1}^{k} (T_a - t_a) \frac{\iota_+^* p_a}{\mathsf{u}}),$$

or equivalently,

$$T_0 = t_0 + \frac{\iota_+^*((T_0' - t_0') + \sum_{a=1}^k (T_a - t_a) p_a)}{\mathsf{u}}. \tag{40}$$

It remains to show that the open and closed mirror maps are given by

$$T_a = t_a + \sum_{i\in I} l_i^{(a)} A_i(q), \quad a = 0, 1, \ldots, k.$$

To see this, we expand the I function from Section **??**:

$$\begin{aligned} I(t, z^{-1}) &= (1 + \frac{t_0' + \sum_{a=1}^k t_a p_a}{z} + O(z^{-2}))(1 + \sum_{\beta>0} q^\beta \prod_{i=1}^r \frac{\prod_{m=-\infty}^0 ((D_i^*)_{\mathbb{T}} + mz)}{\prod_{m=-\infty}^{\langle \beta, D_i\rangle} ((D_i^*)_{\mathbb{T}} + mz)}) \\ &= (1 + \frac{t_0' + \sum_{a=1}^k t_a p_a}{z} + O(z^{-2}))(1 + \sum_{i\in I} z^{-1} A_i(q)(D_i^*)_{\mathbb{T}} + O(z^{-2})). \end{aligned}$$

The z^{-1}-term in $I(t, z^{-1})$ is

$$z^{-1}\text{-term} = z^{-1}((t_0' + \sum_{a=1}^k t_a p_a) + \sum_{i\in I} A_i(q)(D_i^*)_{\mathbb{T}})$$

Therefore, the mirror maps satisfy

$$(T_0' - t_0') + \sum_{a=1}^k (T_a - t_a) p_a = \sum_{i\in I} A_i(q)(D_i^*)_{\mathbb{T}} \tag{41}$$

To obtain the closed mirror map, we consider the nonequivariant version of (41):

$$f_0(q) + \sum_{a=1}^k (T_a - t_a) p_a = \sum_{i\in I} A_i(q) D_i^*$$

where $D_i^* = \sum_{a=1}^k l_i^{(a)} p_a \in H_2(X; \mathbb{Z})$. So $f_0(q) = 0$, and the closed mirror map is given by

$$T_a = t_a + \sum_{i\in I} l_i^{(a)} A_i(q), \quad a = 1, \ldots, k.$$

To obtain the open mirror map, we apply ι_+^* to (41) and restrict to $\mathbb{T}_{L,f}$:

$$\iota_+^* \left((T_0' - t_0') + \sum_{a=1}^k (T_a - t_a) p_a \right) = \sum_{i\in I} A_i(q) \iota_+^* (D_i^*)_{\mathbb{T}_{L,f}} = \sum_{i\in I} A_i(q) l_i^{(0)} \mathsf{u}. \tag{42}$$

Combing (40) and (42), we obtain the open mirror map:

$$T_0 = t_0 + \sum_{i\in I} l_i^{(0)} A_i(q).$$

□

Note that (39) implies

$$Q^\beta = q^\beta \exp\left(\sum_{i\in I} \langle \beta, D_i^* \rangle A_i(q) \right). \tag{43}$$

for all $\beta \in H_2(X; \mathbb{Z})$.

Remark 5.3. The mirror maps (39) are exactly prescribed by the logarithmic solution of the Picard-Fuchs equations (35), as expected.

Corollary 5.4. *Let w be a negative integer, and L be a framed inner brane. Define $l^- = (l_1^-, \dots, l_{k+3}^-)$ to be*

$$l_{i_2}^- = f+n+1, \quad l_{i_3}^- = -(f+n), \quad l_{i_4}^- = -1,$$

and

$$l^- = 0 \text{ if } i \in \{1,2,\dots,k+3\} - \{i_2, i_3, i_4\}.$$

(In other words, $l_i^- = l_i^{(0)} - \langle D_i^*, \alpha\rangle, \quad i = 1, \dots, k+3$.) Then

$$J_w^L(Q;f) = (\frac{q_0/q^\alpha}{Q_0/Q^\alpha})^w \left(1 + \sum_{\beta>0} q^\beta \prod_{i=1}^{k+3} \frac{\prod_{m=-\infty}^{0}(wl_i^- + m)}{\prod_{m=-\infty}^{\langle \beta, D_i^*\rangle}(wl_i^- + m)}\right).$$

Proof. Apply Proposition 5.2 with respect to the fixed point q and L, we have

$$J_w^L(Q;f) = (\frac{q_0^-}{Q_0^-})^w \left(1 + \sum_{\beta>0} q^\beta \prod_{i=1}^{k+3} \frac{\prod_{m=-\infty}^{0}(wl_i^- + m)}{\prod_{m=-\infty}^{\langle \beta, D_i^*\rangle}(wl_i^- + m)}\right),$$

where q_0^- and Q_0^- are open string parameters related by

$$Q_0^- = q_0^- \exp(\sum_{i\in I} l_i^- A_i(q)).$$

Since $l_i^- = l_i^{(0)} - \langle \alpha, D_i^*\rangle$, we have

$$Q_0^- = q_0^- \exp\Big(\sum_{i\in I} l_i^{(0)} A_i(q) - \sum_{i\in I} \langle \alpha, D_i^*\rangle A_i(q)\Big).$$

On the other hand,

$$\frac{Q_0/Q^\alpha}{q_0/q^\alpha} = \exp\Big(\sum_{i\in I} l_i^{(0)} A_i(q) - \sum_{i\in I} \langle \alpha, D_i^*\rangle A_i(q)\Big)$$

and this implies the statement. □

5.3. **Outer brane.** By Corollary 3.6 (a) and Proposition 5.2,

$$\begin{aligned}
F(Q;f) &= \sum_{w>0} q_0^w (-1)^{fw} \frac{\prod_{m=1}^{w-1}(fw+m)}{w\cdot w!} \left(1 + \sum_{\beta>0} q^\beta \prod_{i=1}^{k+3} \frac{\prod_{m=-\infty}^{0}(wl_i^{(0)} + m)}{\prod_{m=-\infty}^{\langle \beta, D_i^*\rangle}(wl_i^{(0)} + m)}\right) \\
&= \sum_{(w,\beta)\in \widetilde{\mathbb{L}}_{\mathrm{eff}}(L), w>0} (-1)^{fw} \frac{\prod_{m=1}^{w-1}(fw+m)}{w\cdot w!} \cdot \frac{1}{\prod_{i\in I\setminus I_0} \langle D_i^*, \beta\rangle!} \\
&\cdot \frac{\prod_{m=-\infty}^{0}(w+m)}{\prod_{m=-\infty}^{\langle D_{i_1}^*, \beta\rangle}(w+m)} \cdot \frac{\prod_{m=-\infty}^{0}(fw+m)}{\prod_{m=-\infty}^{\langle D_{i_2}^*, \beta\rangle}(fw+m)} \cdot \frac{\prod_{m=-\infty}^{0}(-(f+1)w+m)}{\prod_{m=-\infty}^{\langle D_{i_3}^*, \beta\rangle}(-(f+1)w+m)} q_0^w q^\beta \\
&= \sum_{\beta'\in \widetilde{\mathbb{L}}_{\mathrm{eff}}(L)} \frac{(-1)^{\langle D_{i_3}^* + D_{r+3}^*, \beta'\rangle}}{\prod_{i\in \widetilde{I}\setminus\{i_2,i_3\}} \langle D_i^*, \beta'\rangle!} \frac{\prod_{m=-\infty}^{-\langle D_{i_3}^*, \beta'\rangle - 1} m}{\prod_{m=-\infty}^{\langle D_{i_2}^*, \beta'\rangle} m} q^{\beta'}.
\end{aligned}$$

This agrees with Equation (38).

Remark 5.5. The condition $\beta' = (w, \beta) \in \widetilde{\mathbb{L}}_{\mathrm{eff}}(L)$ implies

$$w \neq 0, \quad \langle D_{i_1}^*, \beta'\rangle = w + \sum_{a=1}^{k} d_a l_{i_1}^{(a)} \geq 0.$$

When L is an outer brane, it further implies $w>0$. One may choose appropriate $\mathbb{Z}$-basis $\{l^{(a)}\}_{a=1,\dots,k}$ in $\mathbb{L}$ and $\{p_a\}_{a=1,\dots,k}$ in $\mathbb{L}^\vee$ with respect to the $\mathbb{T}$-fixed point x^+. Since the toric variety $X=(\mathbb{C}^r-Z(\Delta))/G$ is smooth, the G action on $\mathbb{C}^r-Z(\Delta)$ is free; in particular the G action on

$$\{(X_1,\dots,X_r): X_{i_1}=X_{i_2}=X_{i_3}=0, X_i\in\mathbb{C}^* \text{ for other } i\}\subset \mathbb{C}^r-Z(\Delta)$$

is free. It follows that the composition with the projection

$$\mathbb{L}\to \widetilde{N}=\bigoplus_{i=1}^r \mathbb{Z}\widetilde{v}_i\to \bigoplus_{i\in I\backslash I_0}\mathbb{Z}\widetilde{v}_i$$

is an isomorphism. Thus we choose the preimages of $\widetilde{v}_i$, where $i\in I\backslash I_0$ as a basis $\{l^{(a)}\}$ of $\mathbb{L}$. We write $l^{(a_i)}$ to be the preimage of $\widetilde{v}_i$ for $i\in I\backslash I_0$. It is obvious that the cone

$$\sum_{a=1}^k \mathbb{Z}_{\geq 0} l^{(a)}\subset \mathbb{L}_{\text{eff}};$$

while the pairing $\langle p_a, l^{(b)}\rangle=\delta_{ab}$ for the dual basis $p_a\in\mathbb{L}^\vee$, and $p_{a_i}=D_i^*$. With such a choice of $l^{(a)}$ and p_a, the intersection $L\cap X$ is described by the following equations

$$l_{i_1}^{(a)}|X_{i_1}|^2+\sum_{i\in I\backslash I_0} l_i^{(a)}|X_i|^2=r_a,\quad a=1,\dots,k,$$

where $r_1,\dots,r_k$ are Kähler parameters. Since L is an outer brane, if one moves L to infinity, i.e. let $|X_{i_1}|\to\infty$, we will not encounter a $\mathbb{T}$-fixed point, i.e. $|X_i|$ cannot be zero for any $i\in I\backslash I_0$. By our choice of basis $\{l^{(a)}\}$ and $\{p_a\}$ in $\mathbb{L}$ and $\mathbb{L}^\vee$, $l_j^{(a_i)}=\delta_{ij}$ for $i,j\in I\backslash I_0$. Thus $l_{i_1}^{(a)}\leq 0$ for $a=1,\dots,k$, and $\beta'\in\widetilde{\mathbb{L}}_{\text{eff}}(L)$ already implies $w>0$.

This completes the proof of Conjecture 4.3 when L is a framed outer brane.

5.4. **Inner brane.** By Corollary 3.6 (b), Proposition 5.2 and Corollary 5.4,

$$F(Q;f)=I_++I_-,$$

where

$$\begin{aligned}
I_+ &= \sum_{w>0} q_0^w(-1)^{fw}\frac{\prod_{m=1}^{w-1}(fw+m)}{w\cdot w!}\left(1+\sum_{\beta>0}q^\beta\prod_{i=1}^{k+3}\frac{\prod_{m=-\infty}^{0}(wl_i^{(0)}+m)}{\prod_{m=-\infty}^{\langle\beta,D_i^*\rangle}(wl_i^{(0)}+m)}\right)\\
I_- &= \sum_{w<0}(\frac{q_0}{q^\alpha})^w(-1)^{(f+n)w}\frac{\prod_{m=1}^{-w-1}((f+n)(-w)+m)}{(-w)\cdot(-w)!}\left(1+\sum_{\beta>0}q^\beta\prod_{i=1}^{k+3}\frac{\prod_{m=-\infty}^{0}(wl_i^{-}+m)}{\prod_{m=-\infty}^{\langle\beta,D_i^*\rangle}(wl_i^{-}+m)}\right).
\end{aligned}$$

So I_- can be rewritten as

$$\begin{aligned}
I_- &= \sum_{w<0} q_0^w(-1)^{(f+n)w}\frac{\prod_{m=1}^{-w-1}((f+n)(-w)+m)}{(-w)\cdot(-w)!}\left(\sum_{\beta\geq 0}q^{\beta-w\alpha}\prod_{i=1}^{k+3}\frac{\prod_{m=-\infty}^{-w\langle\alpha,D_i^*\rangle}(wl_i^{(0)}+m)}{\prod_{m=-\infty}^{\langle\beta-w\alpha,D_i^*\rangle}(wl_i^{(0)}+m)}\right)\\
&= \sum_{w<0} q_0^w(-1)^{(f+n)w}\frac{\prod_{m=1}^{-w-1}((f+n)(-w)+m)}{(-w)\cdot(-w)!}\left(\sum_{\beta+w\alpha\geq 0}q^{\beta}\prod_{i=1}^{k+3}\frac{\prod_{m=-\infty}^{-w\langle\alpha,D_i^*\rangle}(wl_i^{(0)}+m)}{\prod_{m=-\infty}^{\langle\beta,D_i^*\rangle}(wl_i^{(0)}+m)}\right)
\end{aligned}$$

The calculation of I_+ is identical to the outer brane case.

$$I_+ = \sum_{w>0} q_0^w (-1)^{fw} \frac{\prod_{m=1}^{w-1}(fw+m)}{w\cdot w!} \left(1+\sum_{\beta>0} q^\beta \prod_{i=1}^{k+3} \frac{\prod_{m=-\infty}^{0}(wl_i^{(0)}+m)}{\prod_{m=-\infty}^{\langle \beta, D_i^*\rangle}(wl_i^{(0)}+m)}\right)$$
$$= \sum_{\beta'\in\widetilde{\mathbb{L}}_{\mathrm{eff}}(L), w>0} \frac{(-1)^{\langle D_{i_3}^*+D_{r+3}^*,\beta'\rangle}}{\prod_{i\in\widetilde{I}\setminus\{i_2,i_3\}}\langle D_i^*,\beta'\rangle!} \frac{\prod_{m=-\infty}^{-\langle D_{i_3}^*,\beta'\rangle-1} m}{\prod_{m=-\infty}^{\langle D_{i_2}^*,\beta'\rangle} m} q^{\beta'};$$

while the other half of the contribution I_- is the following

$$I_- = \sum_{\beta'\in\widetilde{\mathbb{L}}_{\mathrm{eff}}(L), w<0} (-1)^{(f+n)w} \frac{\prod_{m=1}^{-w-1}((f+n)(-w)+m)}{(-w)\cdot(-w)!} \cdot \frac{1}{\prod_{i\in I\setminus(I_0\cup\{i_4\})}\langle D_i^*,\beta\rangle!} \cdot \frac{\prod_{m=-\infty}^{-w} m}{\prod_{m=-\infty}^{\langle D_{i_4}^*,\beta\rangle} m}$$
$$\cdot \frac{\prod_{m=-\infty}^{-w}(w+m)}{\prod_{m=-\infty}^{\langle D_{i_1}^*,\beta\rangle}(w+m)} \cdot \frac{\prod_{m=-\infty}^{0}((f+n+1)w+m)}{\prod_{m=-\infty}^{\langle D_{i_2}^*,\beta\rangle}(fw+m)} \cdot \frac{\prod_{m=-\infty}^{0}(-(f+n)w+m)}{\prod_{m=-\infty}^{\langle D_{i_3}^*,\beta\rangle}((-f-1)w+m)} q_0^w \prod_{a=1}^{k} q^\beta$$
$$= \sum_{\beta'\in\widetilde{\mathbb{L}}_{\mathrm{eff}}(L), w<0} \frac{(-1)^{\langle D_{i_3}^*+D_{r+3}^*,\beta'\rangle}}{\prod_{i\in\widetilde{I}\setminus\{i_2,i_3\}}\langle D_i^*,\beta'\rangle!} \frac{\prod_{m=-\infty}^{-\langle D_{i_3}^*,\beta'\rangle-1} m}{\prod_{m=-\infty}^{\langle D_{i_2}^*,\beta'\rangle} m} q^{\beta'}$$

Therefore,

$$F(Q;f) = I_+ + I_- = \sum_{\beta'\in\widetilde{\mathbb{L}}_{\mathrm{eff}}(L)} \frac{(-1)^{\langle D_{i_3}^*+D_{r+3}^*,\beta'\rangle}}{\prod_{i\in\widetilde{I}\setminus\{i_2,i_3\}}\langle D_i^*,\beta'\rangle!} \frac{\prod_{m=-\infty}^{-\langle D_{i_3}^*,\beta'\rangle-1} m}{\prod_{m=-\infty}^{\langle D_{i_2}^*,\beta'\rangle} m} q^{\beta'}.$$

It agrees with Equation (38). This completes the proof of Conjecture 4.3 when L is a framed inner brane.

6. Explicit Mirror Formulae

In this section we list explicit mirror formulae for several examples, whose charge vectors are listed in Table 1. The open and closed mirror maps are given by

$$Q_a = q_a \exp\Big(\sum_{i\in I} l_i^{(a)} A_i(q)\Big), \quad a = 0,\dots,k.$$

The formulae for

$$A_i(q) = \sum_{d_1,\dots,d_k\in\mathbb{Z}} a^i_{d_1,\dots,d_k} \prod_{a=1}^{k} q_a^{d_a}.$$

are listed in Table 2 at the end of this section. In particular, there is no mirror correction when X is the resolved conifold. Notice $A_i(q) = 0$ for $i \neq 1$ for X other than the toric crepant resolution Y_m of $(\mathcal{O}_{\mathbb{P}^1}(-1)\oplus\mathcal{O}_{\mathbb{P}^1}(-1))/\mathbb{Z}_m$.

6.1. $\mathcal{O}_{\mathbb{P}^1}(-1)\oplus\mathcal{O}_{\mathbb{P}^1}(-1)$.

B-model: $\mathbb{T}_q^\vee$ is

$$x_1^{-1}x_2^{-1}x_3x_4 = q = e^t.$$

The mirror curves are listed in the following table.

Phase	Mirror curve
I	$y+1-qy-xy^{-f}=0$
II	$-xy^{-f}+y+1-qxy^{-f+1}=0$

S	$l^{(a)}$, $a = 1, \ldots, k$
$\mathcal{O}_{\mathbb{P}^1}(-1)$	$l^{(1)} = (-1, -1, 1, 1)$
$\mathbb{P}^2$	$l^{(1)} = (-3, 1, 1, 1)$
$\mathbb{P}^1 \times \mathbb{P}^1$	$l^{(1)} = (-2, 1, 1, 0, 0)$ $l^{(2)} = (-2, 0, 0, 1, 1)$
dP_1	$l^{(1)} = (-2, 1, 1, 0, 0)$ $l^{(2)} = (-1, 0, -1, 1, 1)$
dP_2	$l^{(1)} = (-2, 1, 1, 0, 0, 0)$ $l^{(2)} = (-2, 0, 0, 1, 1, 0)$ $l^{(3)} = (-3, 1, 0, 1, 0, 1)$
dP_3	$l^{(1)} = (-2, 1, 1, 0, 0, 0, 0)$ $l^{(2)} = (-2, 0, 0, 1, 1, 0, 0)$ $l^{(3)} = (-3, 1, 0, 1, 0, 1, 0)$ $l^{(4)} = (-3, 0, 1, 0, 1, 0, 1)$
Y_m	$l^{(1)} = (1, 1, 0, -2, 0, 0, \ldots, 0)$ $l^{(2)} = (0, 0, 1, -2, 1, 0, \ldots, 0)$ $l^{(3)} = (0, 0, 0, 1, -2, 1, \ldots, 0)$ $\ldots$ $l^{(m)} = (0, 0, 0, \ldots, 0, 1, -2, 1)$

TABLE 1. charge vectors

A-model:

$$-|X_1|^2 - |X_2|^2 + |X_3|^2 + |X_4|^2 = r,$$

where $r > 0$.

$$|X_4|^2 - |X_2|^2 = c_1, \quad |X_1|^2 - |X_2|^2 = c_2.$$

Therefore

$$(|X_1|^2, |X_2|^2, |X_3|^2, |X_4|^2) = (c_2, 0, r - c_1 + c_2, c_1) + s(1, 1, 1, 1), \quad s \geq 0.$$

See Figure 5.



FIGURE 5. Phase I: $r > c_1 > 0, c_2 = 0$. Phase II: $c_1 = 0, c_2 > 0$.

Mirror Formula: $F(Q; f) = W(q; f) = \sum_{w \neq 0} \sum_{d} n_{d,w}(f) q_0^w q_1^d$.

Phase	(i_1,i_2,i_3)	$n_{d,w}(f)$	$\widetilde{\mathbb{L}}_{\text{eff}}(L)$
I	$(4,1,2)$	$(-1)^{fw+d}\frac{\prod_{m=-d+1}^{d+w-1}(fw+m)}{w\cdot(w+d)!d!}$	$d, w+d\geq 0$
II	$(1,2,4)$	$(-1)^{fw+d}\frac{\prod_{m=-d+1}^{-d+w-1}(fw+m)}{w\cdot(w-d)!d!}$	$w\geq d\geq 0$

6.2. $K_{\mathbb{P}^2}$.

B-model: $\mathbb{T}_q^\vee$ is

$$x_1^{-3}x_2x_3x_4 = q = e^t.$$

The mirror curves are listed in the following table.

Phase	Mirror curve
I	$1-xy^{-f}+y-qx^{-1}y^{f-1}=0$
II	$y+1-xy^{-f}-qx^{-1}y^{f+3}=0$
III	$-xy^{-f}+y+1-qx^3y^{-3f-1}=0$

A-model:

$$|X_2|^2+|X_3|^2+|X_4|^2-3|X_1|^2=r,$$

where $r>0$.

$$|X_2|^2-|X_1|^2=c_1,\quad |X_3|^2-|X_1|^2=c_2.$$

Therefore

$$(|X_1|^2,|X_2|^2,|X_3|^2,|X_4|^2)=(0,c_1,c_2,r-(c_1+c_2))+s(1,1,1,1),\quad s\geq 0.$$

See Figure 6.



FIGURE 6. Phase I: $r>c_1>0, c_2=0$. Phase II: $c_1=0, r>c_2>0$. Phase III: $c_1=c_2<0$.

Mirror Formula: $F(Q;f)=W(q;f)=\sum_{w\neq 0}\sum_d n_{d,w}(f)q_0^w q_1^d$.

Phase	(i_1,i_2,i_3)	$n_{d,w}(f)$	$\widetilde{\mathbb{L}}_{\text{eff}}(L)$
I	$(2,3,1)$	$(-1)^{fw+d}\frac{\prod_{m=d+1}^{3d+w-1}(fw+m)}{w\cdot(w+d)!d!}$	$d, w+d\geq 0$
II	$(3,1,2)$	$(-1)^{fw+d}\frac{\prod_{m=-3d+1}^{-d+w-1}(fw+m)}{w\cdot(w+d)!d!}$	$d, w+d\geq 0$
III	$(1,2,3)$	$(-1)^{fw+d}\frac{\prod_{m=d+1}^{-d+w-1}(fw+m)}{w\cdot(w-3d)!d!}$	$d\geq 0,\ w\geq 3d$

In Phase I, when $f = 0$ we have

$$\sum_{w\neq 0, d\geq 0, w+d\geq 0} N_{0,d\alpha+wb,w}(f) Q_0^w Q_1^d = \sum_{w\neq 0, d\geq 0, w+d\geq 0} (-1)^d \frac{(w+3d-1)!}{w\cdot(w+d)!(d!)^2} q_0^w q_1^d,$$

which agrees with the formula in [16, Section 4.3].

In Phase III, when $f = 0$ we have

$$\sum_{w>0, d\geq 0} N_{0,d\alpha+wb,w}(f) Q_0^w Q_1^d = \sum_{w>0, d\geq 0, w\geq 3d} (-1)^d \frac{(w-d-1)!}{w\cdot(d!)^2(w-3d)!} q_0^w q_1^d,$$

which agrees with [23, Equation A.3].

6.3. $K_{\mathbb{P}^1\times\mathbb{P}^1} = K_{\mathbb{F}_0}$.

B-model: The torus $\mathbb{T}_q^\vee$ is given by the following equations

$$x_1^{-2} x_2 x_3 = q_1 = e^{t_1},$$
$$x_1^{-2} x_4 x_5 = q_2 = e^{t_2}.$$

The mirror curves are listed in the following table.

Phase	Mirror curve
I	$y - q_1 x^{-1} y^{f+2} - x y^{-f} + 1 + q_2 y^2 = 0$
II	$1 + q_1 y^{-1} + y - x y^{-f} - q_2 x^{-1} y^f = 0$
III	$-x y^{-f} + q_1 x^2 y^{-2f} + 1 + y + q_2 x^2 y^{-2f-1} = 0$

A-model:

$$|X_2|^2 + |X_3|^2 - 2|X_1|^2 = r_1, \quad |X_4|^2 + |X_5|^2 - 2|X_1|^2 = r_2$$

where $r_1, r_2 > 0$.

$$|X_3|^2 - |X_1|^2 = c_1, \quad |X_4|^2 - |X_1|^2 = c_2.$$

Therefore

$$(|X_1|^2, |X_2|^2, \ldots, |X_5|^2) = (0, r_1 - c_1, c_1, c_2, r_2 - c_2) + s(1,1,1,1,1), \quad s \geq 0.$$

See Figure 7.



FIGURE 7. Phase I: $r_1 > c_1 > 0, c_2 = 0$. Phase II: $c_1 = 0, r_2 > c_2 > 0$. Phase III: $c_1 = c_2 < 0$

Mirror Formula: $F(Q;f) = W(q;f) = \sum_{w\neq 0}\sum_{d_1,d_2} n_{d_1,d_2,w}(f) q_0^w q_1^{d_1} q_2^{d_2}$.

Phase	(i_1,i_2,i_3)	$n_{d_1,d_2,w}(f)$	$\widetilde{\mathbb{L}}_{\text{eff}}(L)$
I	$(3,1,4)$	$(-1)^{fw+d_2}\frac{\prod_{m=-2d_1-2d_2+1}^{-d_2+w-1}(fw+m)}{w\cdot(w+d_1)!d_1!d_2!}$	$d_1,d_2,w+d_1\geq 0$
II	$(4,3,1)$	$(-1)^{fw}\frac{\prod_{m=d_1+1}^{2d_1+2d_2+w-1}(fw+m)}{w\cdot(w+d_2)!d_1!d_2!}$	$d_1,d_2,w+d_2\geq 0$
III	$(1,4,3)$	$(-1)^{fw+d_1}\frac{\prod_{m=d_2+1}^{-d_1+w-1}(fw+m)}{w\cdot(w-2d_1-2d_2)!d_1!d_2!}$	$d_1,d_2\geq 0,\ w\geq 2d_1+2d_2$

6.4. $K_{dP_1} = K_{\mathbb{F}_1}$.

B-model: The torus $\mathbb{T}_q^\vee$ is given by the following equations

$$x_1^{-2}x_2x_3 = q_1 = e^{t_1},$$
$$x_1^{-1}x_3^{-1}x_4x_5 = q_2 = e^{t_2}.$$

The mirror curves are listed in the following table.

Phase	Mirror curve
I	$y+1+q_1y^2-xy^{-f}-q_1q_2x^{-1}y^{f+3}=0$
II	$1-xy^{-f}-q_1x^{-1}y^f+y-q_1q_2x^{-1}y^{f-1}=0$
III	$1+q_1y^{-1}+y-xy^{-f}-q_2x^{-1}y^{f+1}=0$
IV	$-xy^{-f}+y+q_1x^2y^{-2f-1}+1-q_1q_2x^3y^{-3f-1}=0$
V	$-xy^{-f}+q_1x^2y^{-2f}+1+y-q_2xy^{-f-1}=0$

A-model:

$$|X_2|^2+|X_3|^2-2|X_1|^2 = r_1, \quad |X_4|^2+|X_5|^2-|X_1|^2-|X_3|^2 = r_2$$

where $r_1, r_2 > 0$.

$$|X_4|^2-|X_1|^2 = c_1, \quad |X_2|^2-|X_1|^2 = c_2.$$

Therefore

$$(|X_1|^2,|X_2|^2,\ldots,|X_5|^2) = (0,c_2,r_1-c_2,c_1,r_1+r_2-c_1-c_2)+s(1,1,1,1,1), \quad s\geq 0.$$

See Figure 8.



FIGURE 8. Phase I: $r_1+r_2>c_1>0, c_2=0$. Phase II: $c_1=0, r_1>c_2>0$. Phase III: $r_2>c_1>0, c_2=r_1$. Phase IV: $c_1=c_2<0$. Phase V: $c_1=r_1-c_2<0$.

Mirror Formula: $F(Q;f)=W(q;f)=\sum_{w\neq 0}\sum_{d_1,d_2} n_{d_1,d_2,w}(f)q_0^w q_1^{d_1}q_2^{d_2}$.

Phase	(i_1,i_2,i_3)	$n_{d_1,d_2,w}(f)$	$\tilde{\mathbb{L}}_{\mathrm{eff}}(L)$
I	$(4,1,2)$	$(-1)^{fw+d_1}\frac{\prod_{m=-2d_1-d_2+1}^{-d_1+w-1}(fw+m)}{w\cdot(w+d_2)!(d_1-d_2)!d_2!}$	$w+d_2\geq 0,\ d_1\geq d_2\geq 0$
II	$(2,4,1)$	$(-1)^{fw+d_2}\frac{\prod_{m=d_2+1}^{2d_1+d_2+w-1}(fw+m)}{w\cdot(w+d_1)!(d_1-d_2)!d_2!}$	$w+d_1\geq 0,\ d_1\geq d_2\geq 0$
III	$(4,3,1)$	$(-1)^{fw+d_2}\frac{\prod_{m=d_1-d_2+1}^{2d_1+d_2+w-1}(fw+m)}{w\cdot(w+d_2)!d_1!d_2!}$	$d_1,d_2\geq 0,\ w+d_2\geq 0$
IV	$(1,2,4)$	$(-1)^{fw+d_2}\frac{\prod_{m=d_1+1}^{-d_2+w-1}(fw+m)}{w\cdot(w-2d_1-d_2)!(d_1-d_2)!d_2!}$	$d_1\geq d_2\geq 0,\ w\geq 2d_2+d_2$
V	$(1,4,3)$	$(-1)^{fw+d_1+d_2}\frac{\prod_{m=d_2+1}^{-d_1+d_2+w-1}(fw+m)}{w\cdot(w-2d_1-d_2)!d_1!d_2!}$	$d_1,d_2\geq 0,\ w\geq 2d_1+d_2$

6.5. K_{dP_2}.

B-model: The torus $\mathbb{T}_q^\vee$ is given by

$$x_1^{-2}x_2x_3=q_1=e^{t_1},$$
$$x_1^{-2}x_4x_5=q_2=e^{t_2},$$
$$x_1^{-3}x_2x_4x_6=q_3=e^{t_3}.$$

The mirror curves are listed in the following table.

Phase	Mirror curve
I	$y+1+q_1y^2-q_2x^{-1}y^{f+2}-xy^{-f}-(q_2)^{-1}q_3xy^{-f+1}=0$
II	$1-xy^{-f}+q_1x^2y^{-2f}+q_2y^{-1}+y-(q_2)^{-1}q_3x^{-1}y^{f+1}=0$
III	$1-(q_2)^{-1}q_3xy^{-f-1}-q_1q_2(q_3)^{-1}y^{f+1}-q_2x^{-1}y^f-xy^{-f}+y=0$
IV	$-xy^{-f}+1+q_1x^2y^{-2f}+y+q_2x^2y^{-2f-1}-q_3x^3y^{-3f-1}=0$
V	$-xy^{-f}+y+q_1x^2y^{-2f-1}+q_2x^2y^{-2f}+1-(q_2)^{-1}q_3xy^{-f-1}=0$
VI	$-xy^{-f}-q_2^{-1}q_3xy^{-f+1}-q_1q_2q_3^{-1}xy^{-f-1}+q_2x^2y^{-2f-1}+y+1=0$

A-model:

$$\begin{aligned}|X_2|^2+|X_3|^2-2|X_1|^2 &= r_1,\\ |X_4|^2+|X_5|^2-2|X_1|^2 &= r_2,\\ |X_2|^2+|X_4|^2+|X_6|^2-3|X_1|^2 &= r_3,\end{aligned}$$

where $r_1,r_2,r_3>0$, and $\max(r_1,r_2)<r_3<r_1+r_2$.

$$|X_2|^2-|X_1|^2=c_1,\quad |X_5|^2-|X_1|^2=c_2.$$

Therefore

$$(|X_1|^2,|X_2|^2,\dots,|X_6|^2)=(0,c_1,r_1-c_1,r_2-c_2,c_2,r_3-r_2-c_1+c_2)+t(1,1,1,1,1,1),\quad t\geq 0.$$

See Figure 9.



FIGURE 9. Phase I: $r_2 > c_2 > 0, c_1 = 0$. Phase II: $r_3 - r_2 > c_1 > 0, c_2 = 0$. Phase III: $r_3 - r_2 < c_1 < r_1, c_1 - c_2 = r_3 - r_2$. Phase IV: $r_2 - c_2 = c_1 < 0$ Phase V: $c_1 = c_2 < 0$. Phase VI: $c_1 = r_3 - r_2, c_2 < 0$.

Mirror Formula: $F(Q;f) = W(q;f) = \sum_{w\neq 0} \sum_{d_1,d_2,d_3} n_{d_1,d_2,d_3,w}(f) q_0^w \prod_{a=1}^{3} q_a^{d_a}$.

Phase	(i_1,i_2,i_3)	$n_{d_1,d_2,d_3,w}(f)$	$\widetilde{\mathbb{L}}_{\text{eff}}(L)$
I	$(5,1,2)$	$\frac{(-1)^{fw+d_1+d_3}\prod_{m=-2d_1-2d_2-3d_3+1}^{-d_1-d_3+w-1}(fw+m)}{w(w+d_2)!d_1!d_3!(d_2+d_3)!}$	$d_1,d_3,d_2+d_3 \geq 0$ $w+d_2 \geq 0$
II	$(2,5,1)$	$\frac{(-1)^{fw+d_3}\prod_{m=d_2+1}^{2d_1+2d_2+3d_3+w-1}(fw+m)}{w(w+d_1+d_3)!d_1!d_3!(d_2+d_3)!}$	$d_1,d_3,d_2+d_3 \geq 0$ $w+d_1+d_3 \geq 0$
III	$(5,6,1)$	$\frac{(-1)^{fw+d_3}\prod_{m=d_3+1}^{2d_1+2d_2+3d_3+w-1}(fw+m)}{w(w+d_2)!d_1!(d_1+d_3)!(d_2+d_3)!}$	$d_1,d_1+d_3,d_2+d_3 \geq 0$ $w+d_2 \geq 0$
IV	$(1,4,2)$	$\frac{(-1)^{fw+d_1+d_3}\prod_{m=d_2+d_3+1}^{-d_1-d_3+w-1}(fw+m)}{w(w-2d_1-2d_2-3d_3)!d_1!d_2!d_3!}$	$d_1,d_2,d_3 \geq 0$ $w \geq 2d_1+2d_2+3d_3$
V	$(1,2,5)$	$\frac{(-1)^{fw+d_2}\prod_{m=d_1+d_3+1}^{-d_2+w-1}(fw+m)}{w(w-2d_1-2d_2-3d_3)!d_1!(d_2+d_3)!d_3!}$	$d_1,d_3,d_2+d_3 \geq 0$ $w \geq 2d_1+2d_2+3d_3$
VI	$(1,5,6)$	$\frac{(-1)^{fw+d_3}\prod_{m=d_1+d_3+1}^{-d_3+w-1}(fw+m)}{w(w-2d_1-2d_2-3d_3)!d_1!(d_1+d_3)!(d_2+d_3)!}$	$d_1,d_1+d_3,d_2+d_3 \geq 0$ $w \geq 2d_1+2d_2+3d_3$

6.6. K_{dP_3}.

B-model: The torus $\mathbb{T}_q^\vee$ is given by

$$x_1^{-2}x_2x_3 = q_1 = e^{t_1}, \quad x_1^{-2}x_4x_5 = q_2 = e^{t_2}, \quad x_1^{-3}x_2x_4x_6 = q_3 = e^{t_3}, \quad x_1^{-3}x_3x_5x_7 = q_4 = e^{t_4}.$$

The mirror curves are listed in the following table.

Phase	Mirror curve
I	$1 + y + q_1y^{-1} - q_2x^{-1}y^f - xy^{-f} - (q_2)^{-1}q_3xy^{-f-1} - (q_1)^{-1}q_4x^{-1}y^{f+1} = 0$
II	$-xy^{-f} + 1 + q_1x^2y^{-2f} + q_2x^2y^{-2f-1} + y - (q_2)^{-1}q_3xy^{-f+1} - (q_1)^{-1}q_4xy^{-f-1} = 0$

A-model:

$$\begin{aligned} |X_2|^2+|X_3|^2-2|X_1|^2 &= r_1,\\ |X_4|^2+|X_5|^2-2|X_1|^2 &= r_2,\\ |X_2|^2+|X_4|^2+|X_6|^2-3|X_1|^2 &= r_3,\\ |X_3|^2+|X_5|^2+|X_7|^2-3|X_1|^2 &= r_4, \end{aligned}$$

where $r_1, r_2 > 0$, $\max(r_1,r_2) < r_3 < r_1+r_2$, $\max(r_1,r_2) < r_4 < r_1+r_2$.

$$|X_2|^2-|X_1|^2 = c_1, \quad |X_5|^2-|X_1|^2 = c_2.$$

Therefore

$$\begin{aligned} &(|X_1|^2,|X_2|^2,|X_3|^2,|X_4|^2,|X_5|^2,|X_6|^2,|X_7|^2)\\ =\ &(0,c_1,r_1-c_1,r_2-c_2,c_2,r_3-r_2-c_1+c_2,r_4-r_1+c_1-c_2)+t(1,1,1,1,1,1,1), \quad t\geq 0. \end{aligned}$$

See Figure 10.



FIGURE 10. Phase I: $0 < c_1 < r_4 - r_1$. Phase II: $c_1 = c_2 > 0$.

Mirror Formula: $F(Q;f) = W(q;f) = \sum_{w\neq 0}\sum_{d_1,\dots,d_4} n_{d_1,d_2,d_3,d_4,w}(f) q_0^w \prod_{a=1}^4 q_a^{d_a}$.

Phase	(i_1,i_2,i_3)	$n_{d_1,d_2,d_3,d_4,w}(f)$	$\widetilde{\mathbb{L}}_{\text{eff}}(L)$
I	$(5,2,1)$	$\dfrac{(-1)^{fw+d_3+d_4}\prod_{m=d_1+d_3+1}^{2d_1+2d_2+3d_3+3d_4+w-1}(fw+m)}{w(w+d_2+d_4)!d_3!d_4!(d_2+d_3)!(d_1+d_4)!}$	$d_3,d_4,d_1+d_4,d_2+d_3\geq 0$ $w+d_2+d_4\geq 0$
II	$(1,5,2)$	$\dfrac{(-1)^{fw+d_2+d_4}\prod_{m=d_2+d_4+1}^{-d_1-d_3+w-1}(fw+m)}{w(w-2d_1-2d_2-3d_3-3d_4))!d_3!d_4!(d_2+d_3)!(d_1+d_4)!}$	$d_3,d_4,d_1+d_4,d_2+d_3\geq 0$ $w\geq 2d_1+2d_2+3d_3+3d_4$

6.7. **The toric crepant resolution Y_m of $(\mathcal{O}_{\mathbb{P}^1}(-1)\oplus\mathcal{O}_{\mathbb{P}^1}(-1))/\mathbb{Z}_m$.**

B-model: The torus $\mathbb{T}_q^\vee$ is given by

$$\begin{aligned} x_1x_2x_4^{-2} &= q_1,\\ x_3x_5x_4^{-2} &= q_2,\\ x_4x_6x_5^{-2} &= q_3,\\ &\dots\\ x_{m+1}x_{m+3}x_{m+2}^{-2} &= q_m. \end{aligned}$$

One can write down the equations of the framed mirror curves (33) prescribed in Section 4.1. We do not list them here because they are too long.

A-model:

$$\begin{aligned}
|X_1|^2+|X_2|^2-2|X_4|^2 &= r_1,\\
|X_3|^2+|X_5|^2-2|X_4|^2 &= r_2,\\
|X_4|^2+|X_6|^2-2|X_5|^2 &= r_3,\\
|X_5|^2+|X_7|^2-2|X_6|^2 &= r_4,\\
&\cdots,\\
|X_{m+1}|^2+|X_{m+3}|^2-2|X_{m+2}|^2 &= r_m,
\end{aligned}$$

where $r_1,\ldots,r_m>0$.

The Aganagic-Vafa A-branes in each phase are given as follows:

- I_0:
$$|X_1|^2-|X_4|^2=|X_3|^2-|X_4|^2=-c,\quad c>0$$
- I_b, $1\le b\le m-1$:
$$|X_1|^2-|X_{b+3}|^2=0,\ |X_{b+2}|^2-|X_{b+3}|^2=c,\quad 0<c<r_{b+1}.$$
- I_m:
$$|X_1|^2-|X_{m+3}|^2=0,\ |X_{m+2}|^2-|X_{m+3}|^2=c,\quad c>0.$$
- II_b, $b=1,2$:
$$|X_1|^2-|X_{b+3}|^2=c,\ |X_{b+2}|^2-|X_{b+3}|^2=0,\quad 0<c<r_1.$$
- II_b, $2\le b\le m$:
$$\begin{aligned}&|X_1|^2-|X_{b+3}|^2=c,\ |X_{b+2}|^2-|X_{b+3}|^2=0,\\ &0<c<r_1+2(r_3+2r_4+\cdots+(b-2)r_b).\end{aligned}$$

As an example, the image of the 1-skeleton X_5^1 under the moment map $\mu':X\to(\mathfrak{t}'_{\mathbb{R}})^*$ is in Figure 11.



FIGURE 11. $\mu'(X_5^1)=\mu'((\mathcal{O}(-1)\oplus\mathcal{O}(-1))/\mathbb{Z}_5)^1)$.

Mirror Formula: $F(Q;f)=W(q;f)=\sum_{w\neq 0}\sum_{d_1,\dots,d_m} n_{d_1,\dots,d_m,w}(f)q_0^w\prod_{a=1}^m q_a^{d_a}$.

The mirror formulae for some phases are listed in the following table.

Phase	(i_1,i_2,i_3)	$n_{d_1,\dots,d_m,w}(f)$	$\widetilde{\mathbb{L}}_{\mathrm{eff}}(L)$
I_0	$(4,3,1)$	$(-1)^{fw+d_1}\prod_{m=d_2+1}^{-d_1+w-1}(fw+m)$ $\cdot\frac{1}{w(w-2d_1-2d_2+d_3)!d_1!}$ $\cdot\frac{1}{(d_2-2d_3+d_4)!(d_3-2d_4+d_5)!}$ $\cdots$ $\cdot\frac{1}{(d_{m-2}-2d_{m-1}+d_m)!}\cdot\frac{1}{(d_{m-1}-2d_m)!d_m!}$	$w\geq 2d_1+2d_2-d_3$, $d_1\geq 0$, $d_2-2d_3+d_4\geq 0$, $d_3-2d_4+d_5\geq 0$, $\dots$, $d_{m-2}-2d_{m-1}+d_m\geq 0$, $d_{m-1}-2d_m\geq 0$, $d_m\geq 0$.
I_b $3\leq b\leq m-2$	$(b+2,1,b+3)$	$(-1)^{fw+d_b-2d_{b+1}+d_{b+2}}$ $\cdot\prod_{m=d_1+1}^{-d_b+2d_{b+1}-d_{b+2}+w-1}(fw+m)$ $\cdot\frac{1}{w(w+d_{b-1}-2d_b+d_{b+1})!d_1!d_2!}$ $\cdot\frac{1}{(-2d_1-2d_2+d_3)!(d_3-2d_4+d_5)!}$ $\cdots$ $\cdot\frac{1}{(d_{b-2}-2d_{b-1}+d_b)!(d_{b+1}-2d_{b+2}+d_{b+3})!}$ $\cdots$ $\frac{1}{(d_{m-2}-2d_{m-1}+d_m)!(d_{m-1}-2d_m)!d_m!}$	$w+d_{b-1}-2d_b+d_{b+1}\geq 0$, $d_1,d_2\geq 0$, $-2d_1-2d_2+d_3\geq 0$, $\dots$ $d_{b-2}-2d_{b-1}+d_b\geq 0$, $d_{b+1}-2d_{b+2}+d_{b+3}\geq 0$, $\dots$, $d_{m-2}-2d_{m-1}+d_m\geq 0$, $d_{m-1}-2d_m\geq 0$, $d_m\geq 0$
II_b $3\leq b\leq m-2$	$(1,b+3,b+2)$	$(-1)^{fw+d_{b-1}-2d_b+d_{b+1}}$ $\cdot\prod_{m=d_{b+1}-2d_{b+2}+d_{b+3}+1}^{-d_{b-1}+2d_b-d_{b+1}+w-1}(fw+m)$ $\cdot\frac{1}{w(w+d_1)!d_1!d_2!}$ $\cdot\frac{1}{(-2d_1-2d_2+d_3)!(d_3-2d_4+d_5)!}$ $\cdots$ $\cdot\frac{1}{(d_{b-2}-2d_{b-1}+d_b)!(d_{b+1}-2d_{b+2}+d_{b+3})!}$ $\cdots$ $\cdot\frac{1}{(d_{m-2}-2d_{m-1}+d_m)!(d_{m-1}-2d_m)!d_m!}$	$w+d_1\geq 0$, $d_1,d_2\geq 0$, $-2d_1-2d_2+d_3\geq 0$, $\dots$ $d_{b-2}-2d_{b-1}+d_b\geq 0$, $d_{b+1}-2d_{b+2}+d_{b+3}$, $\dots$, $d_{m-2}-2d_{m-1}+d_m\geq 0$, $d_{m-1}-2d_m\geq 0$, $d_m\geq 0$

S	i	$a^i_{d_1,\dots,d_k}$	constraints on d_a in the sum
$\mathcal{O}_{\mathbb{P}^1}(-1)$	$i=1,\dots,4$	0	
$\mathbb{P}^2$	1	$\frac{(-1)^{d_1-1}(3d_1-1)!}{(d_1!)^3}$	$d_1>0$
	$2,3,4$	0	
$\mathbb{P}^1\times\mathbb{P}^1$	1	$\frac{-(2d_1+2d_2-1)!}{(d_1!)^2(d_2!)^2}$	$d_1,d_2\geq 0$ $(d_1,d_2)\neq(0,0)$
	$2,\cdots,5$	0	
dP_1	1	$\frac{(-1)^{d_2-1}(2d_1+d_2-1)!}{d_1!(d_1-d_2)!(d_2!)^2}$	$d_1\geq d_2\geq 0$ $(d_1,d_2)\neq(0,0)$
	$2,\cdots,5$	0	
dP_2	1	$\frac{(-1)^{d_3-1}(2d_1+2d_2+3d_3-1)!}{d_1!d_2!d_3!(d_1+d_3)!(d_2+d_3)!}$	$d_1,d_2,d_3\geq 0$ $(d_1,d_2,d_3)\neq(0,0,0)$
	$2,\cdots,6$	0	
dP_3	1	$\frac{(-1)^{d_3+d_4-1}(2d_1+2d_2+3d_3+3d_4-1)!}{(d_1+d_3)!(d_1+d_4)!(d_2+d_3)!(d_2+d_4)!d_3!d_4!}$	$d_1+d_3,d_1+d_4\geq 0$ $d_2+d_3,d_2+d_4,d_3,d_4\geq 0$ $(d_1,d_2,d_3,d_4)\neq(0,0,0,0)$
	$2,\cdots,7$	0	
Y_m	$1,2,3$	0	
	4	$\frac{(-1)^{2d_1+2d_2-d_3-1}(2d_1+2d_2-d_3-1)!}{(d_1!)^2d_2!(d_{m-1}-2d_m)!d_m!}\cdot\frac{1}{\prod_{b=2}^{m-2}(d_{b-1}-2d_b+d_{b+1})!}$	each factorial is non-negative $(d_1,\dots,d_m)\neq 0$
	$5,\dots,r-2$	$\frac{(-1)^{d_{i-3}-2d_{i-2}+d_{i-1}-1}}{(d_1!)^2d_2!(-2d_1-2d_2+d_3)!(d_{m-1}-2d_m)!d_m!}\cdot\frac{(-d_{i-3}+2d_{i-2}-d_{i-1}-1)!}{\prod_{b=2,b\neq i-2}^{m-2}(d_{b-1}-2d_b+d_{b+1})!}$	
	$r-1$	$\frac{(-1)^{2d_m-d_{m-1}-1}(2d_m-d_{m-1}-1)!}{(d_1!)^2d_2!(-2d_1-2d_2+d_3)!d_m!}\cdot\frac{1}{\prod_{b=2}^{m-2}(d_{b-1}-2d_b+d_{b+1})!}$	
	r	$\frac{(-1)^{d_m-1}(-d_m-1)!}{(d_1!)^2d_2!(-2d_1-2d_2+d_3)!(d_{m-1}-2d_m)!}\cdot\frac{1}{\prod_{b=2}^{m-2}(d_{b-1}-2d_b+d_{b+1})!}$	

TABLE 2. formulae for $A(q)$ (mirror maps)

References

[1] M. Aganagic, A. Klemm, M. Mariño, and C. Vafa, "The topological vertex," Comm. Math. Phys. **254** (2005), no. 2, 425–478.

[2] M. Aganagic, A. Klemm, C. Vafa, "Disk Instantons, Mirror Symmetry and the Duality Web," Z. Naturforsch. **A 57** (2002), no. 1-2, 128.

[3] M. Aganagic, C. Vafa, "Mirror Symmetry, D-Branes and Counting Holomorphic Discs," arXiv:hep-th/0012041.

[4] V. Bouchard, A. Catuneanu, O. Marchal, and P. Sułlowski, "The remodeling conjecture and the Faber-Pandharipande formula," arXiv:1108.2689.

[5] V. Bouchard, A. Klemm, M. Mariño, and S. Pasquetti, "Remodeling the B-model," Comm. Math. Phys. **287** (2009), no. 1, 117–178.

[6] V. Bouchard, P. Sułkowski, "Topological recursion and mirror curves", arXiv:1105.2052

[7] A. Brini, "Open topological strings and integrable hierarchies: Remodeling the A-model," arXiv:1102.0281.

[8] L. Chen, "Bouchard-Klemm-Mariño-Pasquetti Conjecture for $\mathbb{C}^3$," arXiv:0910.3739.

[9] T. Coates, A. Corti, H. Iritani, H.-H. Tseng, "Quantum cohomology of toric stakcks," in preparation.

[10] B. Eynard and N. Orantin, "Invariants of algebraic curves and topological expansion," Commun. Number Theory Phys. **1** (2007), no. 2, 347–452

[11] K. Fukaya, "Counting pseudo-holomorphic discs in Calabi-Yau 3-fold," arXiv:0908.0148.

[12] W. Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies **131**, Princeton University Press, Princeton, NJ, 1993.

[13] A. Givental, "Equivariant Gromov-Witten invariants", Internat. Math. Res. Notices 1996, no. 13, 613663.

[14] A. Givental, "A mirror theorem for toric complete intersections," Topological field theory, primitive forms and related topics (Kyoto, 1996), 141175, Progr. Math., **160**, Birkhäuser Boston, Boston, MA, 1998.

[15] A. Givental, "Elliptic Gromov-Witten invariants and the generalized mirror conjecture," Integrable systems and algebraic geometry (Kobe/Kyoto, 1997), 107155, World Sci. Publ., River Edge, NJ, 1998.

[16] T. Graber and E. Zaslow, "Open-string Gromov-Witten invariants: calculations and a mirror "theorem"," Orbifolds in mathematics and physics (Madison, WI, 2001), 107–121, Contemp. Math., **310**, Amer. Math. Soc., Providence, RI, 2002.

[17] T. Graber and R. Pandharipande, "Localization of virtual classes," Invent. Math. **135** (1999), no. 2, 487–518.

[18] E.-N. Ionel and T. Parker, "Relative Gromov-Witten invariants," Ann. of Math. (2) **157** (2003), no. 1, 45–96.

[19] E.-N. Ionel and T. Parker, "The symplectic sum formula for Gromov-Witten invariants," Ann. of Math. (2) **159** (2004), no. 3, 935–1025.

[20] S. Katz, C.-C.M. Liu, "Enumerative geometry of stable maps with Lagrangian boundary conditions and multiple covers of the disc," Adv. Theor. Math. Phys. **5** (2001), no. 1, 1–49.

[21] T. Hausel, B. Sturmfels, "Toric hyperKähler varieties," Doc. Math. **7** (2002), 495534.

[22] M. Kontsevich, "Enumeration of rational curves via torus actions," The moduli space of curves (Texel Island, 1994), 335–368, Progr. Math., **129**, Birkhäuser Boston, Boston, MA, 1995.

[23] W. Lerche and P. Mayr, "On $\mathcal{N} = 1$ mirror symmetry for open type II strings," arXiv:hep-th/0111113.

[24] W. Lerche, P. Mayr and Warner, "N=1 special geometry, mixed Hodge variations and toric geometry", arXiv:hep-th/0208039.

[25] A. Li and Y. Ruan, "Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds," Invent. Math. **145** (2001), no. 1, 151–218.

[26] J. Li, "Stable Morphisms to singular schemes and relative stable morphisms," J. Diff. Geom. **57** (2001), 509–578.

[27] J. Li, "A degeneration formula of Gromov-Witten invariants," J. Diff. Geom. **60** (2002), 199-293.

[28] C.-C.M. Liu, K. Liu, and J. Zhou, "A formula of two-partition Hodge integrals," J. Amer. Math. Soc. **20** (2007), no. 1, 149184

[29] J. Li, C.-C.M. Liu, K. Liu, and J. Zhou, "A mathematical theory of the topological vertex," Geom. Topol. **13** (2009), no. 1, 527–621.

[30] B.H. Lian, K. Liu, S.-T. Yau, "Mirror principle I," Asian J. Math. **1** (1997), no. 4, 729–763.

[31] B.H. Lian, K. Liu, S.-T. Yau, "Mirror Principle II," Asian J. Math. **3** (1999), no. 1, 109–146.

[32] B.H. Lian, K. Liu, S.-T. Yau, "Mirror Principle III," Asian J. Math. **3** (1999), no. 4, 771800.

[33] C.-C.M. Liu, "Moduli of J-Holomorphic Curves with Lagrangian Boundary Conditions and Open Gromov-Witten Invariants for an S^1-Equivariant Pair," arXiv:math/0210257.

[34] M. Mariño, "Open string amplitudes and large order behavior in topological string theory," J. High Energy Phys. **2008**, no. 3, 060, 34 pp.

[35] D. Maulik, A. Oblomkov, A. Okounkov, and R. Pandharipande, "Gromov-Witten/Donaldson-Thomas correspondence for toric 3-folds," Invent. Math. **186** (2011), no. 2, 435479.
[36] P. Mayr, "N=1 mirror symmetry and open/closed string duality," Adv. Theor. Math. Phys. **5** (2001), no. 2, 213242.
[37] P. Mayr, "Summing up open string instantons and $N = 1$ string amplitudes", `arXiv:hep-th/0203237`.
[38] J. Solomon, "Intersection theory on the moduli space of holomorphic curves with Lagrangian boundary conditions," `arXiv:math/0606429`.
[39] J.Zhou, "Local Mirror Symmetry for One-Legged Topological Vertex," `arXiv:0910.4320`.
[40] J. Zhou, "Local Mirror Symmetry for the Topological Vertex," `arXiv:0911.2343`.
[41] J. Zhou, "Open string invariants and mirror curve of the resolved conifold," `arXiv:1001.0447`.

Bohan Fang, Department of Mathematics, Columbia University, 2990 Broadway, New York, NY 10027
E-mail address: `b-fang@math.columbia.edu`

Chiu-Chu Melissa Liu, Department of Mathematics, Columbia University, 2990 Broadway, New York, NY 10027
E-mail address: `ccliu@math.columbia.edu`